


## Metadata of the chapter that will be visualized online

Chapter Title	IP <sub>3</sub> Receptor Properties and Function at Membrane Contact Sites		
Copyright Year	2018		
Copyright Holder	Springer International Publishing AG		
Author	Family Name	<b>Roest</b>	
	Particle		
	Given Name	<b>Gemma</b>	
	Suffix		
	Division	Laboratory for Molecular and Cellular Signaling, Department of Cellular and Molecular Medicine & Leuven Kanker Instituut	
	Organization	KU Leuven	
	Address	Leuven, Belgium	
Author	Family Name	<b>Rovere</b>	
	Particle	<b>La</b>	
	Given Name	<b>Rita M.</b>	
	Suffix		
	Division	Laboratory for Molecular and Cellular Signaling, Department of Cellular and Molecular Medicine & Leuven Kanker Instituut	
	Organization	KU Leuven	
	Address	Leuven, Belgium	
Corresponding Author	Family Name	<b>Bultynck</b>	
	Particle		
	Given Name	<b>Geert</b>	
	Suffix		
	Division	Laboratory for Molecular and Cellular Signaling, Department of Cellular and Molecular Medicine & Leuven Kanker Instituut	
	Organization	KU Leuven	
	Address	Leuven, Belgium	
	Email	geert.bultynck@med.kuleuven.be 	
Corresponding Author	Family Name	<b>Parys</b>	
	Particle		
	Given Name	<b>Jan B.</b>	
	Suffix		
	Division	Laboratory for Molecular and Cellular Signaling, Department of Cellular and Molecular Medicine & Leuven Kanker Instituut	

Organization KU Leuven

Address Leuven, Belgium

Email jan.parys@med.kuleuven.be



---

Abstract

The inositol 1,4,5-trisphosphate (IP<sub>3</sub>) receptor (IP<sub>3</sub>R) is a ubiquitously expressed Ca<sup>2+</sup>-release channel localized in the endoplasmic reticulum (ER). The intracellular Ca<sup>2+</sup> signals originating from the activation of the IP<sub>3</sub>R regulate multiple cellular processes including the control of cell death versus cell survival via their action on apoptosis and autophagy. The exact role of the IP<sub>3</sub>Rs in these two processes does not only depend on their activity, which is modulated by the cytosolic composition (Ca<sup>2+</sup>, ATP, redox status, . . .) and by various types of regulatory proteins, including kinases and phosphatases as well as by a number of oncogenes and tumor suppressors, but also on their intracellular localization, especially at the ER-mitochondrial and ER-lysosomal interfaces. At these interfaces, Ca<sup>2+</sup> microdomains are formed, in which the Ca<sup>2+</sup> concentration is finely regulated by the different ER, mitochondrial and lysosomal Ca<sup>2+</sup> transport systems and also depends on the functional and structural interactions existing between them. In this review, we therefore discuss the most recent insights in the role of Ca<sup>2+</sup> signaling in general and of the IP<sub>3</sub>R in particular in the control of basal mitochondrial bioenergetics, apoptosis, and autophagy at the level of inter-organellar contact sites.

---

Keywords

(separated by '-')

Apoptosis - Autophagy - Ca<sup>2+</sup> microdomains - Cell death - Cell survival - Endoplasmic reticulum - IP<sub>3</sub> receptor - Lysosomes - Membrane contact sites - Mitochondria

---

## Chapter 7

# IP<sub>3</sub> Receptor Properties and Function at Membrane Contact Sites

Gemma Roest, Rita M. La Rovere, Geert Bultynck, and Jan B. Parys

**Abstract** The inositol 1,4,5-trisphosphate (IP<sub>3</sub>) receptor (IP<sub>3</sub>R) is a ubiquitously expressed Ca<sup>2+</sup>-release channel localized in the endoplasmic reticulum (ER). The intracellular Ca<sup>2+</sup> signals originating from the activation of the IP<sub>3</sub>R regulate multiple cellular processes including the control of cell death versus cell survival via their action on apoptosis and autophagy. The exact role of the IP<sub>3</sub>Rs in these two processes does not only depend on their activity, which is modulated by the cytosolic composition (Ca<sup>2+</sup>, ATP, redox status, ...) and by various types of regulatory proteins, including kinases and phosphatases as well as by a number of oncogenes and tumor suppressors, but also on their intracellular localization, especially at the ER-mitochondrial and ER-lysosomal interfaces. At these interfaces, Ca<sup>2+</sup> microdomains are formed, in which the Ca<sup>2+</sup> concentration is finely regulated by the different ER, mitochondrial and lysosomal Ca<sup>2+</sup> transport systems and also depends on the functional and structural interactions existing between them. In this review, we therefore discuss the most recent insights in the role of Ca<sup>2+</sup> signaling in general and of the IP<sub>3</sub>R in particular in the control of basal mitochondrial bioenergetics, apoptosis, and autophagy at the level of inter-organellar contact sites.

**Keywords** Apoptosis · Autophagy · Ca<sup>2+</sup> microdomains · Cell death · Cell survival · Endoplasmic reticulum · IP<sub>3</sub> receptor · Lysosomes · Membrane contact sites · Mitochondria

Gemma Roest and Rita M. La Rovere are Joint first authors.

AU1

G. Roest · R. M. La Rovere · G. Bultynck (✉) · J. B. Parys (✉)  
Laboratory for Molecular and Cellular Signaling, Department of Cellular and Molecular  
Medicine & Leuven Kanker Instituut, KU Leuven, Leuven, Belgium  
e-mail: [geert.bultynck@med.kuleuven.be](mailto:geert.bultynck@med.kuleuven.be); [jan.parys@med.kuleuven.be](mailto:jan.parys@med.kuleuven.be)



© Springer International Publishing AG 2018

J. Krebs (ed.), *Membrane Dynamics and Calcium Signaling*, Advances in  
Experimental Medicine and Biology 981,

[https://doi.org/10.1007/978-3-319-55858-5\\_7](https://doi.org/10.1007/978-3-319-55858-5_7)

## 24 Abbreviations

25	ALS	Amyotrophic lateral sclerosis
26	AMPK	AMP-activated kinase
27	ATG	Autophagy-related
28	BIRD-2	Bcl-2/IP <sub>3</sub> R disruptor-2 peptide
29	CICR	Ca <sup>2+</sup> -induced Ca <sup>2+</sup> release
30	CREB	cAMP response element-binding protein
31	DT40 TKO	DT40 IP <sub>3</sub> R triple knock-out
32	ER	Endoplasmic reticulum
33	Fis1	Fission 1 homologue
34	GRP75	Glucose-regulated protein 75
35	GRP78/BiP	Glucose-regulated protein 78
36	GSK3 $\beta$	Glycogen synthase kinase-3 $\beta$
37	IMM	Inner mitochondrial membrane
38	IBC	IP <sub>3</sub> -binding core
39	IP <sub>3</sub>	Inositol 1,4,5-trisphosphate
40	IP <sub>3</sub> R	IP <sub>3</sub> receptor
41	LC3	Microtubule-associated protein light chain 3
42	LRRK2	Leucine-rich repeat kinase 2
43	MAM	Mitochondria-associated ER membrane
44	MCU	Mitochondrial Ca <sup>2+</sup> uniporter
45	Mfn	Mitofusin
46	mPTP	Mitochondrial permeabilization transition pore
47	mTORC1	Mechanistic target of rapamycin complex 1
48	OMM	Outer mitochondrial membrane
49	NAADP	Nicotinic acid adenine dinucleotide phosphate
50	PACS-2	Phosphofurin acidic cluster sorting protein 2
51	PERK	Protein kinase RNA-like ER kinase
52	PIP <sub>3</sub>	Phosphatidylinositol 3,4,5-trisphosphate
53	PKB/Akt	Protein kinase B
54	PML	Promyelocytic leukemia
55	PTEN	Phosphatase and tensin homolog
56	PTPIP51	Protein tyrosine phosphatase-interacting protein-51
57	ROS	Reactive oxygen species
58	RyR	Ryanodine receptor
59	SERCA	Sarco-/endoplasmic reticulum Ca <sup>2+</sup> ATPase
60	TCA	Tricarboxylic acid
61	TFEB	Transcription factor EB
62	TMX	Thioredoxin-like transmembrane protein
63	TPC	Two-pore channel
64	TRPML	Transient receptor potential mucolipin
65	ULK1/2	Atg1/Unc-51-like kinase 1/2
66	UPR	Unfolded protein response
67	VAPB	Vesicle-associated protein B
68	VDAC	Voltage-dependent anion channel


## 7.1 The IP<sub>3</sub> Receptor, the Main Ca<sup>2+</sup>-Release Channel of the Endoplasmic Reticulum

69

Complex spatio-temporal Ca<sup>2+</sup> signals regulate many fundamental cellular processes including fertilization, differentiation, proliferation, gene transcription, metabolism, contraction, secretion, etc. [1, 2]. The endoplasmic reticulum (ER) is the main Ca<sup>2+</sup>-storage organelle and therefore plays a central role in intracellular Ca<sup>2+</sup> signaling. ER Ca<sup>2+</sup> handling depends on three major mechanisms, Ca<sup>2+</sup> uptake via the sarco-/endoplasmic Ca<sup>2+</sup>-ATPases (SERCA) [3], Ca<sup>2+</sup> storage by various luminal Ca<sup>2+</sup>-binding proteins [4] and controlled Ca<sup>2+</sup> release. Besides passive Ca<sup>2+</sup> leak through still largely unidentified basal ER Ca<sup>2+</sup>-leak channels, Ca<sup>2+</sup> release out of the ER primarily occurs via the inositol 1,4,5-trisphosphate (IP<sub>3</sub>) receptor (IP<sub>3</sub>R) and the ryanodine receptor (RyR). While the RyR is predominantly expressed in a limited number of tissues, especially muscles and brain [5], the IP<sub>3</sub>R is ubiquitously expressed and present in virtually every cell type [6]. IP<sub>3</sub>Rs are activated by IP<sub>3</sub> produced by phospholipase C after cell activation, e.g. by extracellular agonists, hormones, growth factors or neurotransmitters, and play a crucial role in the initiation and propagation of intracellular and intercellular Ca<sup>2+</sup> signals [7–10].

In *Mammalia* and other higher organisms, three different genes encode IP<sub>3</sub>Rs (*ITPR1*, *ITPR2* and *ITPR3*) giving rise to three isoforms named IP<sub>3</sub>R1, IP<sub>3</sub>R2, and IP<sub>3</sub>R3 that display at the amino acid level an overall similarity of 75–80% [11]. Each of these isoforms is about 2700 amino acid long and assembles into tetrameric structures with a total molecular mass of 1.2 MDa. Splice isoforms and the possibility to form both homo- and heterotetramers increase the diversity between IP<sub>3</sub>Rs. Each monomeric isoform has the same general structure containing five distinct functional domains (Fig. 7.1): the N-terminal coupling domain usually called the suppressor domain (for IP<sub>3</sub>R1: a.a. 1–225), the IP<sub>3</sub>-binding core (IBC, a.a. 226–578) containing the IP<sub>3</sub>-binding site, the central coupling domain or modulatory and transducing domain (a.a. 579–2275), the channel domain containing six transmembrane helices (a.a. 2276–2589) and finally the C-terminal tail also named gatekeeper domain (a.a. 2590–2749) [13, 14].

At the structural level progress has followed two main paths. First, X-ray crystallography permitted to obtain high-resolution structural information of the N-terminal part of the IP<sub>3</sub>R, i.e. the suppressor domain and the IBC [15, 16]. The latter contains two domains, a first one containing a  $\beta$ -trefoil fold (IBC- $\beta$ ) and a second one containing an armadillo repeat fold (IBC- $\alpha$ ) with the cleft between them forming the IP<sub>3</sub>-binding site. Their structure was also resolved at high resolution (3–3.8 Å) both for the apo- and the IP<sub>3</sub>-bound form, elucidating changes occurring during IP<sub>3</sub>R activation [17, 18]. In the presence of IP<sub>3</sub>, a rearrangement of IBC- $\beta$  and IBC- $\alpha$  occurs, constraining the IP<sub>3</sub>-binding cleft. Additionally, the suppressor domain, originally in contact with both IBC- $\beta$  and IBC- $\alpha$ , re-orientates itself, probably to couple to the channel domain for its activation. Recently, Mikoshiba and co-workers succeeded in obtaining the crystal structure for the 2217 a.a. long cytosolic part of IP<sub>3</sub>R1 at a resolution of 5.8–7.4 Å [19]. Comparison of the structures of



**Fig. 7.1** Linear representation of the type 1 IP<sub>3</sub>R isoform and its regulation by proteins involved in the control of cell survival and cell death. The five distinct IP<sub>3</sub>R1 functional domains are delineated: the suppressor domain, the IP<sub>3</sub>-binding core, the modulatory and transducing domain, the channel domain containing the six trans-membrane helices (indicated as black bars) and the gatekeeper domain. Oncogenes are represented in green, while tumor suppressors are indicated in red. Bar-headed lines indicate an inhibitory interaction and arrow-headed lines indicate a stimulatory interaction. For more information, see text. Reproduced with modifications from own previous work ([12], <https://doi.org/10.1016/j.bbamcr.2016.01.002>)

different truncated variants obtained in the absence and the presence of IP<sub>3</sub> supports a long-range gating mechanism in which the signal is transferred from the IBC via two  $\alpha$ -helical domains (HD1, a.a. 605–1009 and HD3, a.a. 1593–2217) to a so-called leaflet (corresponding to a.a. 2195–2215 of HD3) that relays the IP<sub>3</sub>R conformational change to the channel pore. Strikingly, the HD3/leaflet region was essential for the IP<sub>3</sub>R Ca<sup>2+</sup>-release activity in intact cells exposed to extracellular agonists.

Second, cryo-electron microscopy allowed gaining an increasingly better view of the full-length IP<sub>3</sub>R structure. Serysheva and co-workers obtained by single-particle cryo-electron microscopy about 85% of the structure of rat cerebellar IP<sub>3</sub>R1 at a 4.7 Å resolution in the apo-state [20, 21], confirming the close apposition of the N- and C-termini.

This structure enables the binding of four IP<sub>3</sub> molecules to a tetrameric IP<sub>3</sub>R, but the exact stoichiometry underlying IP<sub>3</sub>R opening by IP<sub>3</sub> remained for a long time elusive. The elegant work of Yule and co-workers, based on the ectopic expression in null-background HEK293 cells of concatenated IP<sub>3</sub>Rs of which the IP<sub>3</sub>-binding sites could be mutated, indicated that IP<sub>3</sub>R-mediated Ca<sup>2+</sup> release is only triggered when 4 IP<sub>3</sub> molecules are bound to the tetrameric IP<sub>3</sub>R [22]. This stringent condition will not only prevent spurious IP<sub>3</sub>R opening, but also enables the buffering of a significant amount of IP<sub>3</sub> molecules by the IP<sub>3</sub>R channels themselves, ensuring that sufficiently high IP<sub>3</sub> concentrations are reached before IP<sub>3</sub>R opening takes place [23].

Besides the physiological agonist IP<sub>3</sub>, two metabolites produced by *Penicillium brevicompactum*, the adenophostins A and B, can activate the IP<sub>3</sub>R at a much lower concentration than IP<sub>3</sub> itself [24]. Cytosolic Ca<sup>2+</sup> is considered an important co-agonist of the IP<sub>3</sub>R that acts in a bell-shaped dependent manner: a low [Ca<sup>2+</sup>] (typically ≤0.3 μM) enhances IP<sub>3</sub>-induced Ca<sup>2+</sup> release, while a high [Ca<sup>2+</sup>] (above 0.3 μM) suppresses IP<sub>3</sub>-induced Ca<sup>2+</sup> release [25–28]. ATP [29] and oxidizing conditions as e.g. thimerosal [30, 31] were also shown to stimulate IP<sub>3</sub>-induced Ca<sup>2+</sup> release. Finally, cAMP has been reported to sensitize IP<sub>3</sub>Rs [32] independently of any phosphorylation event [33]. Moreover, very recently evidence was presented that when cAMP is produced, IP<sub>3</sub> can induce Ca<sup>2+</sup> release from a different Ca<sup>2+</sup> store that is not sensitive to IP<sub>3</sub> alone [34].

Interestingly, each of the IP<sub>3</sub>R isoforms acts as a signaling hub, able to integrate various cellular inputs and to deliver a specific output signal that elicits a specific cellular response [9, 10, 35]. It should therefore be emphasized that although all IP<sub>3</sub>R isoforms function along the same basic mechanisms, important differences in sensitivity exists between the various IP<sub>3</sub>R isoforms, with IP<sub>3</sub>R2 being the most atypical one [36]. The sensitivity of the IP<sub>3</sub>R isoforms to IP<sub>3</sub> follows the rank-order IP<sub>3</sub>R2 > IP<sub>3</sub>R1 > IP<sub>3</sub>R3 [37–41]. In addition, the sensitivity to cytosolic Ca<sup>2+</sup> varies between the various IP<sub>3</sub>R isoforms [42–44]. ATP similarly has a biphasic effect on IP<sub>3</sub>R activity with between the IP<sub>3</sub>R isoforms different sensitivities as well for the stimulatory [40, 45, 46] as for the inhibitory phase [47]. Moreover, IP<sub>3</sub>R2, the isoform that is most sensitive to stimulatory ATP concentrations [45], proved to be dominant in a heterotetrameric receptor [48]. Finally, the oxidizing agent thimerosal has a biphasic action on IP<sub>3</sub>R1 but only an inhibitory one on IP<sub>3</sub>R3 [46, 49] while cAMP seems to have its main (or only) effect on IP<sub>3</sub>R2 [32].

Inhibition of IP<sub>3</sub>R activity is generally achieved by one of the following compounds: heparin, 2-APB, xestospongin C (or B) and caffeine. Except for the problem that all those compounds display non-specific effects [50], an additional problem is that they also may display a differential sensitivity between the various IP<sub>3</sub>R isoforms [51]. The synthesis of IP<sub>3</sub> derivatives may however provide in the future more specific and/or more potent inhibitors [52].

Further regulation of IP<sub>3</sub>R activity can be obtained by various post-translational modifications, including palmitoylation [53], cross-linking by the action of transglutaminase 2 [54] and phosphorylation/dephosphorylation events triggered by multiple protein kinases and protein phosphatases [55, 56]. Finally, the existence of multiple regulatory or scaffolding proteins able to associate with the IP<sub>3</sub>R tetramer and to modulate its activity should be emphasized. Over 100 proteins have already been shown to interact with and regulate the IP<sub>3</sub>R [57], including IRBIT [58, 59], calmodulin [60–64], neuronal CaBP1 [65–67], the anti-apoptotic Bcl-2 [68–70], Bcl-XL [71] and reticulocalbin 1 [72], the pro-autophagic Beclin 1 [73, 74], the chaperones ERp44 [75], GRP78/BiP [76] and sigma-1 receptor [77] and cytochrome c [78].

Many of these associated proteins, including several oncogenes and tumor suppressors (Fig. 7.1), are involved in the regulation of apoptosis and autophagy [12, 79]. This correlates with the important function of IP<sub>3</sub>Rs and IP<sub>3</sub>R-mediated Ca<sup>2+</sup> signals in the control of cell-fate decisions, including processes related to cell death and survival

177 [80]. More specifically, the role of the IP<sub>3</sub>R and of IP<sub>3</sub>-induced Ca<sup>2+</sup> release has been  
 178 recognized in both apoptosis [81–86] and autophagy [85–90]. These processes depend  
 179 on the activity of the IP<sub>3</sub>R and its regulation by associated proteins and protein kinases,  
 180 as well as on its intracellular localization, especially at the interface between ER and  
 181 mitochondria or at the interface between ER and lysosomes.

## 182 **7.2 Mitochondrial and Lysosomal Ca<sup>2+</sup> Handling in Cell** 183 **Death and Survival Processes**

### 184 **7.2.1 Apoptosis and Its Regulation by Ca<sup>2+</sup>**

185 Apoptosis is the major form of programmed cell death and can be initiated by either  
 186 an extrinsic or an intrinsic –mitochondrial- pathway that both converge at the level of  
 187 the activation of the effector caspases (caspase-3, -6 and -7) [91–93].

188 In the intrinsic pathway, Ca<sup>2+</sup> overload of the mitochondrial matrix is a well-known  
 189 factor leading to apoptosis [94]. Interestingly, several links exist between Ca<sup>2+</sup> and  
 190 apoptosis. In a first mechanism, mitochondrial Ca<sup>2+</sup> is thought to bind cardiolipin, which  
 191 thereby dissociates from mitochondrial complex II. The mitochondrial complex II  
 192 disintegrates, leading to the release of the succinate dehydrogenase A and B subunits.  
 193 This results in a massive reactive oxygen species (ROS) production and opening of the  
 194 mitochondrial permeability transition pore (mPTP) [95]. This mPTP was proposed to  
 195 arise from dimers of the F-ATP synthase in the inner mitochondrial membrane (IMM)  
 196 [96, 97], though there still exists some controversy on the exact subunits involved  
 197 [98]. Recent data indicate that mPTP opening can occur subsequently to Ca<sup>2+</sup> binding  
 198 to the  $\beta$  subunit of the catalytic F1 part [99]. This likely results in a conformational  
 199 change transmitted via the oligomycin-sensitivity conferring protein subunit to the  
 200 peripheral stalk and the IMM.

201 The mPTP is a high conductance channel that upon its opening results in the  
 202 collapse of the electrochemical potential and in a loss of ion and solute distribution,  
 203 terminating ATP synthesis. Subsequently to mPTP opening, solutes enter the mito-  
 204 chondrial matrix, leading to mitochondrial swelling, rupture of the outer mito-  
 205 chondrial membrane (OMM) and release of the pro-apoptotic factors located in the  
 206 cristae, including cytochrome c. The release of these factors to the cytoplasm in  
 207 turn leads to caspase activation and apoptosis [100].

208 Besides mPTP opening, Ca<sup>2+</sup> can also indirectly activate mitochondrial apopto-  
 209 sis. Cytosolic Ca<sup>2+</sup> can activate calcineurin, which dephosphorylates the sensitizer  
 210 BH3-only protein Bad. This will cause its release from 14-3-3 proteins and allow its  
 211 interaction with anti-apoptotic Bcl-XL. The latter will then be unable to counteract  
 212 Bax/Bak activation and the subsequent permeabilization of the OMM will lead to  
 213 apoptosis [101].

214 Finally, prolonged, severe depletion of the ER Ca<sup>2+</sup> store will lead to ER  
 215 stress and activation of the unfolded protein response (UPR) to re-establish ER



homeostasis. If the latter cannot be achieved, the UPR will eventually promote the expression of pro-apoptotic proteins, including members of the Bcl-2 family like BH3-only proteins, as well as that of GADD34 and of ERO1 $\alpha$ , all leading to the cell demise by apoptosis [102, 103].

## 7.2.2 Autophagy and Its Regulation by Ca<sup>2+</sup>

Autophagy is a highly conserved degradation pathway in which cellular components are targeted to the lysosomes. Basal and stress-induced autophagy act as survival pathways ensuring that protein aggregates, long-lived proteins, lipids, dysfunctional organelles or intracellular pathogens are digested and their constituents recycled, contributing to protein quality control, energy and cellular homeostasis [104]. Yet, excessive autophagy, brought about by the autophagy-inducing Tat-Beclin 1 peptide or by a severe/on-going nutrient starvation stress, can result in autophagic cell death (or so-called “autosis”) via a process that is regulated by the Na<sup>+</sup>/K<sup>+</sup>-ATPase pump [105, 106].

Macroautophagy, generally referred to as autophagy, is the best-studied type of autophagy. It is characterized by the de novo formation of double-membrane vesicles named autophagosomes that engulf cytosolic proteins and/or organelles, the “cargo”. These autophagosomes eventually fuse with the lysosome to form autolysosomes in which the cargo is degraded.

Main upstream controllers of autophagy are AMP-activated kinase (AMPK) and mechanistic target of rapamycin complex 1 (mTORC1). AMPK triggers autophagy in two ways, as it can phosphorylate and inhibit mTORC1, while directly activating the Atg1/Unc-51-like kinase 1/2 (ULK1/2) complex. The process downstream of ULK1/2 is further controlled by many proteins including about 30 autophagy-related (Atg) proteins [107, 108].

The initiation of the autophagy can start at various intracellular compartments, though the ER –and the ER-mitochondrial contact sites– seems to form a preferential location [109–111]. The original structure forming on the ER has been named the omegasome, which subsequently expands to form the phagophore [112]. Its subsequent elongation and closure, mediated by the Atg12-Atg5 complex, and the lipidation of microtubule-associated protein light chain 3 (LC3) by Atg4 and Atg8, lead to the formation of the autophagosomes. These phagosomes will eventually fuse either directly, or indirectly via prior fusion with an endosome, with lysosomes, forming hereby the autolysosomes needed for cargo degradation [107, 113].

The regulation of autophagy by Ca<sup>2+</sup> appears quite complex as intracellular Ca<sup>2+</sup> can stimulate or even be essential for autophagy [54, 73, 114–118] while it also can inhibit it [119–122]. Possible explanations to resolve the apparent controversy include the use of various cell types and various autophagy induction methods, and the fact that Ca<sup>2+</sup> can act on multiple check points upstream of autophagy as well as during the autophagic process [85, 87–89, 123–127]. In that respect, the intracellular localization of the IP<sub>3</sub>Rs appears important. Ca<sup>2+</sup> signals directed to the

mitochondria will e.g. control mitochondrial metabolism and thus impact autophagy differently than  $\text{Ca}^{2+}$  signals directed to other targets. The importance for cell survival/cell death of  $\text{IP}_3\text{R}$  localization at membrane contact sites will therefore be discussed in the subsequent sections.


### 7.3 The Role of the $\text{IP}_3\text{R}$ at ER: Mitochondrial Contact Sites

$\text{Ca}^{2+}$  signals originating from the ER critically control cell death and survival by impacting the mitochondria [94, 128, 129]. This is possible through the existence of sites of close apposition between the ER and the mitochondria, covering about 10% of the mitochondrial surface, depending on the metabolic state of the cells [130]. These sites are based on the existence of structural connections between both organelles, resulting in the formation of mitochondria-associated ER membranes (MAMs). In those MAMs, numerous proteins were identified that play a role in  $\text{Ca}^{2+}$  handling, lipid transfer, inter-organelle tethering, regulation of mitochondrial fission and fusion, cell death control and mitochondrial metabolism [131–136].

The distance between both organelles is an important element in the control of cell death versus cell survival [134]. The contact sites are dynamic structures in which the distance between both membranes as well as the length of these contact sites vary in function of the physiological conditions. In general, the contact sites run over several hundreds of nanometers of length and the distance between smooth ER and mitochondria range between 10 and 50 nm and between rough ER and mitochondria between 50 and 80 nm [130]. By using synthetic linkers, ER-mitochondrial distance could experimentally be fourfold reduced increasing simultaneously fourfold the length of the ER-mitochondrial interface [137]. This resulted in an accelerated mitochondrial  $\text{Ca}^{2+}$  accumulation and an increased  $\text{Ca}^{2+}$ -dependent activation of the mPTP. Such tightening of the contact between ER and mitochondria physiologically occurs e.g. in cells undergoing apoptosis.

In physiological conditions, the distance between ER and mitochondria is determined by proteins responsible for tethering both organelles together. Although the molecular identity of the tethers remains less well understood in higher organisms than in yeast, several proteins contributing to ER-mitochondrial tethering have already been identified [133, 134, 138] (Fig. 7.2).

In the first place, one should mention that the ER-localized  $\text{IP}_3\text{Rs}$  are in the MAMs physically coupled to the voltage-dependent anion channels (VDAC) 1 located in the OMM via chaperones like glucose-regulated protein 75 (GRP75) [140] (Fig. 7.2a). This tethering enables an efficient “quasi-synaptic”  $\text{Ca}^{2+}$  transfer between ER and mitochondria [141]. This structure leads to high local  $[\text{Ca}^{2+}]$  (in the range of 10–20  $\mu\text{M}$ ) in the ER-mitochondrial interspace [142, 143], which overcomes the low-affinity properties of the mitochondrial  $\text{Ca}^{2+}$  uniporter (MCU) of the IMM [144]. Through the existence of these  $\text{Ca}^{2+}$  microdomains, the mitochondria



**Fig. 7.2** The ER-mitochondrial tethers. Schematic visualization of the most important ER-mitochondrial tethers: (a) the IP<sub>3</sub>R-GRP75-VDAC tether, (b) the mitofusin-based tethering, (c) the VAPB-PTPIP51 tether and (d) the Bap31-Fis1 tether. For more information, see text. Reproduced from ([139], <https://doi.org/10.1016/j.tins.2016.01.008>) available through <https://creativecommons.org/licenses/by/4.0/>

can be regulated even by low-level Ca<sup>2+</sup> signals. Dependently on the Ca<sup>2+</sup> signal properties either cell survival or cell death can be promoted.

Mitofusin (Mfn) 2, enriched in the MAMs, was proposed to form homo- and heterodimers with mitochondria-localized Mfn2 and Mfn1 respectively (Fig. 7.2b), thereby establishing proper ER-mitochondrial apposition [145, 146]. Mfn2 function can be modulated by various proteins, including trichoplein/mitostatin, which counteracts Mfn2 function [147], and the ubiquitin ligase MITOL, which stimulates Mfn2 degradation [148]. In agreement with the model that Mfn2 has a tethering function, both trichoplein/mitostatin and MITOL impair mitochondrial Ca<sup>2+</sup> uptake and protect against Ca<sup>2+</sup>-dependent apoptosis. Other studies however have challenged the concept of Mfn2 as an ER-mitochondrial tether and actually proposed that it functions as an anti-tethering protein. This was based on the observation that cells lacking Mfn2 display an increased ER-mitochondrial tethering and an augmented agonist-induced mitochondrial Ca<sup>2+</sup> uptake [149–152]. Resolving this controversy will probably depend on obtaining more detailed insights in the molecular composition of the ER-mitochondrial interface [153, 154].

Other proteins endowed with a role as tethers between the ER and the mitochondria are the scaffolds based on (a) vesicle-associated protein B (VAPB) in the ER interacting with protein tyrosine phosphatase-interacting protein-51 (PTPIP51) of the OMM, (b) on Bap31 in the ER and Fission 1 homologue (Fis1) in the OMM and (c) the ER stress sensor protein kinase RNA-like ER kinase (PERK) [133, 134, 138].

VABP is an integral ER membrane protein which mutation occurs in forms of motor neuron diseases including amyotrophic lateral sclerosis (ALS) [155]. Its interaction via its N-terminus with PTPIP51 (Fig. 7.2c) controls ER-mitochondria coupling and  $\text{Ca}^{2+}$  transfer between both organelles [156]. Moreover, the VAPB P56S mutant, which causes familial type 8 ALS [157], demonstrates both an increased binding to PTPIP51 and an increased  $\text{Ca}^{2+}$  flux to the mitochondria.


A pro-apoptotic interaction of Bap31 with Fis1 has been demonstrated (Fig. 7.2d), though the role of this interaction under normal physiological conditions is not yet understood [158]. The pro-apoptotic effect is due to the fact that this platform can both recruit pro-caspase-8 and increase  $\text{Ca}^{2+}$  transfer from the ER to the mitochondria. Moreover, the multifunctional phosphofurin acidic cluster sorting protein 2 (PACS-2) can modulate ER-mitochondria coupling and this has been related to Bap31 cleavage [159, 160].

Finally, PERK contributes to ER-mitochondria tethering independently of its canonical role in the UPR pathway, since in PERK-deficient cells the expression of a kinase-dead PERK was sufficient to re-establish ER-mitochondrial contacts and to re-sensitize them to apoptosis [161].

### 7.3.1 *IP<sub>3</sub>R-Mediated $\text{Ca}^{2+}$ Signals in Cell Survival*

Basal IP<sub>3</sub>R-mediated  $\text{Ca}^{2+}$  signals usually in the form of  $\text{Ca}^{2+}$  oscillations- drive mitochondrial bioenergetics (Fig. 7.3a), especially by stimulating the activity of pyruvate dehydrogenase, isocitrate dehydrogenase and  $\alpha$ -ketoglutarate dehydrogenase. These enzymes participate in the tricarboxylic acid (TCA) cycle and so control adequate ATP synthesis [162, 163]. In addition to this, mitochondrial  $\text{Ca}^{2+}$  can also enhance the activity of the ATP synthase complex V and of the adenine nucleotide translocator [164]. Furthermore, IP<sub>3</sub>R-driven  $\text{Ca}^{2+}$  oscillations also sustain the mitochondrial metabolism by driving the transcription of the gene for MCU via regulation of the transcription factor CREB [165].

DT40 cells devoid of all IP<sub>3</sub>R isoforms (DT40 TKO, ~~developed by~~ [166]) demonstrated in comparison to their wild-type counterparts a very different basic energy metabolism, with an increased Warburg effect and increased ROS production, explaining their reduced proliferation [167]. These effects may be aggravated by reduced MCU expression in cells lacking IP<sub>3</sub>R expression and thus  $\text{Ca}^{2+}$  oscillations [165]. Inhibition of IP<sub>3</sub>Rs in non-tumor cells led to a decreased ATP production, a subsequent activation of AMPK and therefore an increase in pro-survival basal autophagy [168]. The latter effect might depend on growth conditions as the increase in AMPK activity in DT40 TKO cells was observed in one study [168] but not in another [167]. Interestingly, similar results were obtained when ER-mitochondria coupling was impaired by downregulation of VABP or of PTPIP51 [169]. Tightening of the ER-mitochondria connection by a synthetic linker or by overexpression of either VABP or PTPIP51 reversed the effect. However, overexpression of the tether proteins only impaired autophagy induced by mTORC1



**Fig. 7.3**  $\text{Ca}^{2+}$  transfer between ER and mitochondria and between ER and lysosomes. Schematic representation of  $\text{Ca}^{2+}$  handling in juxta-ER microdomains. (a)  $\text{Ca}^{2+}$  transfer to the mitochondria is mediated by the close connection between the IP<sub>3</sub>R and in the ER and VDAC in the outer mitochondrial membrane (OMM). Moreover, additional tethers (the mitofusin tether is depicted) contribute to the close apposition of ER and mitochondria. The  $\text{Ca}^{2+}$  in the mitochondrial matrix drives mitochondrial bio-energetics via stimulation of the TCA cycle, the ATP synthase complex V and the adenine nucleotide translocator. In case of mitochondrial  $\text{Ca}^{2+}$  overload, the mPTP is opened, causing mitochondrial swelling and apoptosis. (b) Regulation of the  $\text{Ca}^{2+}$  concentration in the microdomain between ER and lysosomes involves both the IP<sub>3</sub>R in the ER membrane and the lysosomal  $\text{Ca}^{2+}$  channels as TRPML1 and TPC2. Lysosomal  $\text{Ca}^{2+}$  uptake likely occurs via a  $\text{Ca}^{2+}/\text{H}^{+}$  exchanger and molecular identification of ER-lysosomal tethering is presently lacking. The  $\text{Ca}^{2+}$  in the microdomain between ER and lysosomes can regulate the autophagic process upstream of autophagosome formation, during autophagic flux and via calcineurin-mediated activation of transcription factor EB.  $\text{Ca}^{2+}$  fluxes are depicted by red arrows. The green arrows indicate stimulatory interactions. For more information, see text.

inhibition and not nutrient starvation-induced autophagy, uncovering differences
between the role of mitochondrial  $\text{Ca}^{2+}$  handling in different types of autophagy.

In contrast, the survival of tumorigenic cells depends on the IP<sub>3</sub>R<sub>s</sub> as adequate  $\text{Ca}^{2+}$ 
signaling is necessary to sustain mitochondrial metabolism [170, 171]. Although
tumor cells proliferate in an uncontrolled way independently of the mitochondrial
bioenergetics, they need the TCA cycle for the production of metabolites as nucle-
otides for cell growth and proliferation. The AMPK activation and the subsequent
autophagy occurring after IP<sub>3</sub>R inhibition is therefore not sufficient to sustain cell
survival, resulting in a mitotic catastrophe.

Of note, the net effect of IP<sub>3</sub>Rs on mitochondrial bio-energetics will depend on the local activity of the SERCA pumps at the MAMs (Fig. 7.3a), which, by pumping Ca<sup>2+</sup> back into the ER, will limit the available Ca<sup>2+</sup> in this microdomain and thus counteract ER-mitochondrial Ca<sup>2+</sup> transfer. The local SERCA activity can be dynamically regulated, for instance by associated proteins like thioredoxin-like transmembrane protein (TMX) 1 [172, 173]. This redox-sensitive oxidoreductase is enriched at the MAMs, locally inhibiting SERCA pump activity and thus causing a local increase in [Ca<sup>2+</sup>] at the ER-mitochondrial interface and enhancing oxidative phosphorylation. In contrast, low levels of TMX1 will enhance SERCA activity, shifting Ca<sup>2+</sup> away from the mitochondria and from the Ca<sup>2+</sup>-driven bio-energetic pathway. In fact, loss of TMX1 has been observed in tumor cells, favoring their growth by potentially contributing to the Warburg effect. Nevertheless, a low-level ER-mitochondrial Ca<sup>2+</sup> transfer must remain present to sustain proper TCA cycling and the production of the mitochondrial intermediates necessary for anabolic processes that ensure cell survival upon cell division [80, 170].

Apart from a regulation of autophagy, mitochondrial Ca<sup>2+</sup> also impacts mitophagy, the specific elimination of mitochondria via the autophagic process. In fibroblasts expressing a mutation in the electron transport chain leading to a mild phenotype, mitochondrial Ca<sup>2+</sup> uptake decreased, leading to increased autophagy, increased mitophagy and increased biogenesis of new mitochondria [174]. This adaptive response may account for the limited phenotype observed in these cells. In human RPE1 cells, ER-mitochondrial contacts appeared necessary for starvation-induced autophagy and for mitophagy [175]. The sigma-1 receptor located at the MAMs was crucial for starvation-induced autophagy. Its downregulation, however, did not affect mitophagy, and neither did the downregulation of all the RyR isoforms, of all the SERCA isoforms, or of any of the three IP<sub>3</sub>R isoforms. Since the combined knockdown of all three IP<sub>3</sub>R isoforms or of the MCU inhibited mitophagy, this specifically indicates the importance of IP<sub>3</sub>R-mediated Ca<sup>2+</sup> transfer from the ER to the mitochondria. Noteworthy, the sigma-1 receptor at the MAMs stabilizes IP<sub>3</sub>R3 and so regulates Ca<sup>2+</sup> transfer between ER and mitochondria [176] (Fig. 7.3a).

The anti-apoptotic proteins Bcl-2, Bcl-XL and Mcl-1, of which at least the first two have been identified in the MAM fraction [136] (Fig. 7.3a), can bind to the gatekeeper domain in the C-terminal region of the IP<sub>3</sub>R (Fig. 7.1), which is activated by its interaction with the N-terminal IP<sub>3</sub>-binding domain [177]. Their binding leads to a sensitization of the IP<sub>3</sub>R, increased transfer of Ca<sup>2+</sup> to the mitochondria and therefore an increased production of ATP, promoting cell survival [69, 71, 178]. Recent work uncovered the existence of a biphasic action of Bcl-XL on the IP<sub>3</sub>R, i.e. a high-affinity activation at low Bcl-XL concentration and a lower affinity inhibition at high Bcl-XL concentrations [179]. Finally, it should be mentioned that Bcl-XL bound to the IP<sub>3</sub>R C-terminus in the presence of active K-Ras4B phosphorylated by protein kinase C, a trimolecular complex is formed that antagonizes the anti-apoptotic action of Bcl-XL and promotes cell death via excessive autophagy [180].



### 7.3.2 IP<sub>3</sub>R-Mediated Ca<sup>2+</sup> Signals in Apoptosis

412

In contrast, excessive IP<sub>3</sub>R-mediated Ca<sup>2+</sup> release directed to the mitochondria will lead to mitochondrial Ca<sup>2+</sup> overload and eventual apoptosis, as discussed above (Sect. 7.2.1). Especially IP<sub>3</sub>R3 [181] and VDAC1 [182] were proposed to play an important role in pro-apoptotic Ca<sup>2+</sup> signaling, but depending on the cell type, also other IP<sub>3</sub>R isoforms can participate in such pro-apoptotic Ca<sup>2+</sup> signaling [183, 184]. Moreover, besides IP<sub>3</sub>Rs, other Ca<sup>2+</sup>-transport systems that can impact ER-to-mitochondria Ca<sup>2+</sup> flux can be enriched in the MAMs, including SERCA pumps [185] and S1T, a truncated form of SERCA1, induced upon ER stress and unable to pump Ca<sup>2+</sup> [186]. By forming an ER Ca<sup>2+</sup>-leak pathway, S1T expression leads to ER depletion, mitochondrial immobilization and increased ER-mitochondrial contact sites leading to increased mitochondrial Ca<sup>2+</sup> levels and apoptosis.

At the level of the MAMs, multiple proteins, including oncogenes and tumor suppressors, regulate IP<sub>3</sub>R activity and thus ER-mitochondria Ca<sup>2+</sup> transfer and impact the apoptosis process [12].

Various anti-apoptotic Bcl-2 family members (Fig. 7.3a), including Bcl-2, Bcl-XL and Mcl-1, can modulate IP<sub>3</sub>R activity [12, 187–191]. Bcl-2 binds to all three IP<sub>3</sub>R isoforms at a conserved stretch of 20 amino acids located in the middle of their modulatory and transducing domain (Fig. 7.1), inhibiting IP<sub>3</sub>R-mediated Ca<sup>2+</sup> release [192]. Bcl-2 in this way prevents pro-apoptotic Ca<sup>2+</sup> signaling. The BH4 domain of Bcl-2 is hereby sufficient to inhibit IP<sub>3</sub>Rs and to protect cells against Ca<sup>2+</sup>-dependent apoptosis [70]. Of note, the BH4 domain of Bcl-2 appears to be unique compared to the related domain present in Bcl-XL, which was much less effective in binding to and inhibiting IP<sub>3</sub>Rs [193]. In contrast, the BH4 domain of Bcl-XL appeared more effective for suppressing VDAC1-mediated pro-apoptotic Ca<sup>2+</sup> transfer to the mitochondria [194].

Consequently, disrupting this interaction by using a peptide corresponding to the binding site of Bcl-2 on IP<sub>3</sub>R1 resulted in increased IP<sub>3</sub>R activity and apoptosis [192]. Based on this binding site, the cell-permeable Bcl-2/IP<sub>3</sub>R disruptor-2 peptide (BIRD-2) was developed [195]. It elicited spontaneous pro-apoptotic Ca<sup>2+</sup>-release events in various types of cancer cells, including chronic lymphocytic leukemia [195], diffuse large B-cell lymphoma [183], multiple myeloma and follicular lymphoma [196] and small cell lung cancer [197].

IP<sub>3</sub>R activity can also be reduced by phosphorylation by the pro-survival protein kinase B (PKB/Akt) thereby preventing pro-apoptotic Ca<sup>2+</sup> release [198–200]. The PKB/Akt phosphorylation site is conserved among all three IP<sub>3</sub>R isoforms (Fig. 7.1), but a more selective role for PKB/Akt in the regulation of IP<sub>3</sub>R3 was proposed, as a study demonstrated that PKB/Akt could only dampen the Ca<sup>2+</sup> signals in cells expressing that isoform [201]. The phosphatase and tensin homolog (PTEN), a phosphatidylinositol 3,4,5-trisphosphate (PIP<sub>3</sub>) phosphatase upstream of PKB/Akt, counteracts the effect of PKB/Akt on IP<sub>3</sub>Rs while re-expression of PTEN in PTEN-deficient cancer cells restored their sensitivity to pro-apoptotic Ca<sup>2+</sup>-dependent stimuli [200]. Further studies indicated a direct role for PTEN on

IP<sub>3</sub>R phosphorylation and function at the MAMs, where PTEN functioned as a protein phosphatase counteracting PKB/Akt-mediated phosphorylation of IP<sub>3</sub>R channels and augmenting ER-mitochondrial Ca<sup>2+</sup> flux [202]. In addition to this, it was very recently discovered that PTEN promotes Ca<sup>2+</sup>-induced apoptosis in a phosphatase-independent way [203]. Indeed, PTEN competes with the F-box protein FBXL2, an ubiquitin E3 ligase component belonging to the SCF (SKP1, Cullin 1, F-box protein) E3 ligase family [204] with known tumor suppressor action by arresting the cell cycle [205, 206], for binding to the IP<sub>3</sub>R channels. FBXL2 is responsible for IP<sub>3</sub>R ubiquitination and its subsequent proteasomal degradation. As a consequence, PTEN, by competing with FBXL2, stabilizes IP<sub>3</sub>R and promotes continuous Ca<sup>2+</sup> transfer to the mitochondria leading to apoptosis. Finally, the negative regulation of IP<sub>3</sub>R by PKB/Akt is antagonized by the tumor suppressor promyelocytic leukemia (PML) that recruits protein phosphatase PP2A to the MAMs. The latter dephosphorylates PKB/Akt, inhibiting its kinase activity and thus suppressing PKB/Akt-mediated phosphorylation of IP<sub>3</sub>R (Fig. 7.1). Through this mechanism, PML increases Ca<sup>2+</sup> release by the IP<sub>3</sub>R, supporting mitochondrial Ca<sup>2+</sup> overload and apoptosis [207].

In the heart, it was demonstrated that cyclophilin D participates to the IP<sub>3</sub>R-GRP75-VDAC complex and regulates Ca<sup>2+</sup> transfer to the mitochondria [208]. Its genetic or pharmacological inhibition attenuated mitochondrial Ca<sup>2+</sup> overload and protected the cells from the consequences of hypoxia-reoxygenation.

Glycogen synthase kinase-3β (GSK3β) is another regulator of the IP<sub>3</sub>R located in the MAMs [209]. GSK3β phosphorylates the IP<sub>3</sub>R and increases IP<sub>3</sub>-induced Ca<sup>2+</sup> release. Upon ischemia-reperfusion injury, GSK3β activity is increased in cardiomyocytes, leading to increased pro-apoptotic Ca<sup>2+</sup> signaling towards the mitochondria and cell death. Conversely, pharmacological inhibition of GSK3β protects the heart against damage subsequent to ischemia-reperfusion injury.

Finally, it should also be mentioned that increased transfer of Ca<sup>2+</sup> to the mitochondria can also lead to cellular senescence. Adequate Ca<sup>2+</sup> transfer to the mitochondria via IP<sub>3</sub>R2 and the MCU appeared important in oncogene-induced senescence and replicative senescence [210].

## 7.4 The Role of the IP<sub>3</sub>R at ER: Lysosomal Contact Sites

Lysosomes do not only play a fundamental role in the degradation of proteins and organelles, but are also bona fide Ca<sup>2+</sup> stores [211–213] containing Ca<sup>2+</sup> at a concentration of about 500 μM while their pH is between 4 and 5 [214]. Ca<sup>2+</sup> uptake in the lysosomes is usually assumed to be mediated by a Ca<sup>2+</sup>/H<sup>+</sup> exchanger [215] though other possibilities exist. In contrast to the uncertainty concerning the uptake mechanism, multiple Ca<sup>2+</sup>-release channels are known to be present in the lysosomes, including transient receptor potential mucolipin (TRPML) 1 and two-pore channel (TPC) 2 [216]. Moreover, the lysosomes were identified as a target for nicotinic acid adenine dinucleotide phosphate (NAADP), a very potent agent eliciting Ca<sup>2+</sup> release even at nanomolar concentrations (for a recent review, please



see [217]). With respect to the Ca<sup>2+</sup>-releasing agent NAADP, most evidence point towards channels of the TPC family as their target but other channels, including TRPML1, have also been proposed. Further work is needed to fully understand the function of the various TPC channels, as work by various groups have shown that they can permeate several ions, including Ca<sup>2+</sup>, Na<sup>+</sup> and H<sup>+</sup>, and that their properties can be strongly dependent on experimental conditions and/or the presence of regulatory factors [218, 219].

As the autophagy process is regulated by Ca<sup>2+</sup> (see above, Sect. 7.2.2) and as the Ca<sup>2+</sup> ion has only a short range of messenger action [220], it can be assumed that the Ca<sup>2+</sup> concentration in the microdomain between the ER and the lysosomes plays an important role in the control and/or fine-tuning of the autophagic process. This concentration will depend on the relative activity of the Ca<sup>2+</sup> transporters in ER and lysosomes, their mutual functional interaction and the relative distance between the two organelles [85] (Fig. 7.3b).

In that respect, it is important to note that the two main Ca<sup>2+</sup> channels expressed in the lysosomes, i.e. TRPML1 [221–223] and TPC2 [219, 224, 225] have already been implicated in autophagy (see further below).

As is the case for the mitochondria, also a very close association between ER and lysosomes was documented. Extensive contact sites between the two organelles were visualized through electron microscopy, showing in primary human fibroblasts that over 80% of the lysosomes are forming contacts of <20 nm with the ER [226]. This close connection should allow for mutual functional interactions between ER and lysosomes. In rat pulmonary artery smooth muscle cells, the close contacts between lysosomes and the sarcoplasmic reticulum allow for the functional coupling of lysosomes to RyR3 via Ca<sup>2+</sup>-induced Ca<sup>2+</sup> release (CICR) [227]. The contact sites involved show a membrane separation of about 15 nm apart, and extend for over 300 nm; increasing the membrane separation to 50 nm leads to a failure in activating RyR3. In contrast to what is known about other contact sites, the molecular mechanisms responsible for ER-lysosomal tethering are still only very partially recognized [228, 229].

Several studies have demonstrated that a small amount of Ca<sup>2+</sup> released by NAADP can trigger much larger Ca<sup>2+</sup> signals by activation of IP<sub>3</sub>Rs or RyRs located on the ER by CICR or by ER Ca<sup>2+</sup> store overload [226, 230–232] (Fig. 7.3b). The converse situation is also possible and Ca<sup>2+</sup> released by the ER can affect lysosomal Ca<sup>2+</sup> handling by stimulation of either NAADP synthesis or of the NAADP-induced Ca<sup>2+</sup> release [233]. The Ca<sup>2+</sup> ions released by the IP<sub>3</sub>R can also be taken up in the lysosomes, leading to a dampening of the Ca<sup>2+</sup> signals [234, 235]. A similar functional interaction also exist between active TRPML1 channel and ER Ca<sup>2+</sup> release [236].

Taken together, this indicates that Ca<sup>2+</sup> microdomains exist between the ER and the lysosomes in which the local Ca<sup>2+</sup> concentration is regulated in a very complex way, whereby the IP<sub>3</sub>R plays an important role [85, 213, 237–240] (Fig. 7.3b). These Ca<sup>2+</sup> microdomains can therefore regulate the autophagy process. Evidence for stimulatory [241, 242] and inhibitory [243, 244] effects of lysosomal Ca<sup>2+</sup> release on autophagy were presented. These differences in effects can be due to the use of different cellular models and of different mechanisms of autophagy induction differentially affecting the

relation between ER and lysosomes, but can also be an indication that  $\text{Ca}^{2+}$  can impact in a different way the various stages of the autophagic pathway. Moreover, the various lysosomal  $\text{Ca}^{2+}$  channels have been linked to different parts of the autophagy process. The absence of TRPML1 led to an upregulation of autophagy induction but an impaired lysosomal fusion resulting in the accumulation of significantly larger vesicles with a higher  $[\text{Ca}^{2+}]$  [245]. TRPML1 was upregulated during nutrient starvation [246] and is important for the centripetal movement of lysosomes after autophagy induction [247]. In cardiomyocytes, TPCs seem essential for both basal and induced autophagy [248]. Moreover, TPCs can be regulated by the leucine-rich repeat kinase 2 (LRRK2). The latter is a multifunctional protein, of which various mutated species have been implicated in various diseases and Parkinson's disease in particular. In the cell, it participates in the control of various important processes including autophagy [249]. It now appears that LRRK2 stimulates NAADP-induced  $\text{Ca}^{2+}$  release from the lysosomes [241] (Fig. 7.3b). Furthermore, the aberrant lysosomal morphology observed in fibroblasts from patients harboring the LRRK2 G2019S mutation—the most prominent one in Parkinson's disease—can be rescued by TPC2 inhibition or knockout [250]. The local role of  $\text{Ca}^{2+}$  in this process is underpinned by the fact that BAPTA-AM, a fast intracellular  $\text{Ca}^{2+}$  buffer, but not EGTA-AM, a slow intracellular  $\text{Ca}^{2+}$  buffer, could also rescue the aberrant lysosomal morphology.

Finally, nutrient starvation induced TRPML1-mediated  $\text{Ca}^{2+}$  release out of the lysosomes, locally activating calcineurin [251]. Calcineurin subsequently dephosphorylated transcription factor EB (TFEB) leading to its nuclear translocation. Importantly, TFEB is a master regulator of many genes of the lysosomal/autophagic pathway [252], including the gene for TRPML1 itself [253, 254]. This demonstrates that TRPML can mediate its own upregulation through  $\text{Ca}^{2+}$ -dependent activation of TFEB. Moreover, this upregulation appears essential for adapting lysosomes to conditions of nutrient starvation [246].

These results indicate that the  $\text{IP}_3\text{R}$ , together with the lysosomal  $\text{Ca}^{2+}$  channels, participate in the occurrence of local  $\text{Ca}^{2+}$  signals in the microdomain between ER and lysosomes and can thus impact autophagy progression at various levels.

## 7.5 Conclusions

The  $\text{IP}_3\text{R}$ , by evoking intracellular  $\text{Ca}^{2+}$  signals with a specific spatio-temporal profile, is well known to regulate multiple cellular processes, ranging from fertilization to cell death via control of differentiation/proliferation, metabolism, contraction, secretion and many other processes. The  $\text{IP}_3\text{R}$ , of which three isoforms exist, is finely regulated by a variety of cytosolic factors including  $\text{Ca}^{2+}$  itself and ATP, as well as by a multitude of regulatory proteins, including cell death and survival proteins and kinases and phosphatases. The intracellular localization of the  $\text{IP}_3\text{R}$  at ER-mitochondrial and at ER-lysosomal contact sites is however also very important for its fundamental role in regulating cellular bioenergetics, autophagy and apoptosis. The existence of  $\text{Ca}^{2+}$  microdomains at these inter-organellar interfaces and the possibility to locally control  $[\text{Ca}^{2+}]$  in these microdomains, distinct from the bulk

cytosolic [Ca<sup>2+</sup>], allows for a versatile role of Ca<sup>2+</sup> signaling in a cell's decision to engage cell death, cell survival or cell adaptation processes in basal and stress-related conditions.

**Acknowledgements** GR is recipient of a Ph.D. fellowship of the Research Fund—Flanders (FWO). Work performed in the laboratory of the authors was supported by research grants of the FWO, the Research Council of the KU Leuven and the Interuniversity Attraction Poles Programmes (Belgian Science Policy).

## References

- Berridge MJ, Lipp P, Bootman MD (2000) The versatility and universality of calcium signalling. *Nat Rev Mol Cell Biol* 1:11–21. <https://doi.org/10.1038/35036035>
- Berridge MJ, Bootman MD, Roderick HL (2003) Calcium signalling: dynamics, homeostasis and remodelling. *Nat Rev Mol Cell Biol* 4:517–529. <https://doi.org/10.1038/nrm1155>
- Vandecaetsbeek I, Vangheluwe P, Raeymaekers L et al (2011) The Ca<sup>2+</sup> pumps of the endoplasmic reticulum and Golgi apparatus. *Cold Spring Harb Perspect Biol* 3:a004184. <https://doi.org/10.1101/cshperspect.a004184>
- Prins D, Michalak M (2011) Organellar calcium buffers. *Cold Spring Harb Perspect Biol* 3:a004069. <https://doi.org/10.1101/cshperspect.a004069>
- Lanner JT (2012) Ryanodine receptor physiology and its role in disease. *Adv Exp Med Biol* 740:217–234. [https://doi.org/10.1007/978-94-007-2888-2\\_9](https://doi.org/10.1007/978-94-007-2888-2_9)
- Vermassen E, Parys JB, Mauger J-P (2004) Subcellular distribution of the inositol 1,4,5-trisphosphate receptors: functional relevance and molecular determinants. *Biol Cell* 96:3–17. <https://doi.org/10.1016/j.biocel.2003.11.004>
- Fedorenko OA, Popugaeva E, Enomoto M et al (2014) Intracellular calcium channels: inositol-1,4,5-trisphosphate receptors. *Eur J Pharmacol* 739:39–48. <https://doi.org/10.1016/j.ejphar.2013.10.074>
- Foskett JK, White C, Cheung K-H, Mak D-OD (2007) Inositol trisphosphate receptor Ca<sup>2+</sup> release channels. *Physiol Rev* 87:593–658. <https://doi.org/10.1152/physrev.00035.2006>
- Mikoshiha K (2015) Role of IP<sub>3</sub> receptor signaling in cell functions and diseases. *Adv Biol Regul* 57:217–227. <https://doi.org/10.1016/j.jbior.2014.10.001>
- Parys JB, De Smedt H (2012) Inositol 1,4,5-trisphosphate and its receptors. *Adv Exp Med Biol* 740:255–279. [https://doi.org/10.1007/978-94-007-2888-2\\_11](https://doi.org/10.1007/978-94-007-2888-2_11)
- Taylor CW, Genazzani AA, Morris SA (1999) Expression of inositol trisphosphate receptors. *Cell Calcium* 26:237–251. <https://doi.org/10.1054/ceca.1999.0090>
- Bittremieux M, Parys JB, Pinton P, Bultynck G (2016) ER functions of oncogenes and tumor suppressors: Modulators of intracellular Ca<sup>2+</sup> signaling. *Biochim Biophys Acta* 1863:1364–1378. <https://doi.org/10.1016/j.bbamcr.2016.01.002>
- Bosanac I, Michikawa T, Mikoshiha K, Ikura M (2004) Structural insights into the regulatory mechanism of IP<sub>3</sub> receptor. *Biochim Biophys Acta* 1742:89–102. <https://doi.org/10.1016/j.bbamcr.2004.09.016>
- Uchida K, Miyauchi H, Furuichi T et al (2003) Critical regions for activation gating of the inositol 1,4,5-trisphosphate receptor. *J Biol Chem* 278:16551–16560. <https://doi.org/10.1074/jbc.M300646200>
- Bosanac I, Alattia J-R, Mal TK et al (2002) Structure of the inositol 1,4,5-trisphosphate receptor binding core in complex with its ligand. *Nature* 420:696–700. <https://doi.org/10.1038/nature01268>

16. Bosanac I, Yamazaki H, Matsu-Ura T et al (2005) Crystal structure of the ligand binding suppressor domain of type 1 inositol 1,4,5-trisphosphate receptor. *Mol Cell* 17:193–203. <https://doi.org/10.1016/j.molcel.2004.11.047>
17. Lin C-C, Baek K, Lu Z (2011) Apo and InsP<sub>3</sub>-bound crystal structures of the ligand-binding domain of an InsP<sub>3</sub> receptor. *Nat Struct Mol Biol* 18:1172–1174. <https://doi.org/10.1038/nsmb.2112>
18. Seo M-D, Velamakanni S, Ishiyama N et al (2012) Structural and functional conservation of key domains in InsP<sub>3</sub> and ryanodine receptors. *Nature* 483:108–112. <https://doi.org/10.1038/nature10751>
19. Hamada K, Miyatake H, Terauchi A, Mikoshiba K (2017) IP<sub>3</sub>-mediated gating mechanism of the IP<sub>3</sub> receptor revealed by mutagenesis and X-ray crystallography. *Proc Natl Acad Sci USA* 114:4661–4666. <https://doi.org/10.1073/pnas.1701420114>
20. Fan G, Baker ML, Wang Z et al (2015) Gating machinery of InsP<sub>3</sub>R channels revealed by electron cryomicroscopy. *Nature* 527:336–341. <https://doi.org/10.1038/nature15249>
21. Serysheva II, Baer M, Fan G (2017) Structural insights into IP<sub>3</sub>R function. *Adv Exp Med Biol* 2017; (this issue).
22. Alzayady KJ, Wang L, Chandrasekhar R et al (2016) Defining the stoichiometry of inositol 1,4,5-trisphosphate binding required to initiate Ca<sup>2+</sup> release. *Sci Signal* 9:ra35. <https://doi.org/10.1126/scisignal.aad6281>
23. Taylor CW, Konieczny V (2016) IP<sub>3</sub> receptors: take four IP<sub>3</sub> to open. *Sci Signal* 9:pe1. <https://doi.org/10.1126/scisignal.aaf6029>
24. Takahashi M, Kagasaki T, Hosoya T, Takahashi S (1993) Adenophostins A and B: potent agonists of inositol-1,4,5-trisphosphate receptor produced by *Penicillium brevicompactum*. Taxonomy, fermentation, isolation, physico-chemical and biological properties. *J Antibiot (Tokyo)* 46:1643–1647
25. Bezprozvanny I, Watras J, Ehrlich BE (1991) Bell-shaped calcium-response curves of Ins (1,4,5)P<sub>3</sub>- and calcium-gated channels from endoplasmic reticulum of cerebellum. *Nature* 351:751–754. <https://doi.org/10.1038/351751a0>
26. Finch EA, Turner TJ, Goldin SM (1991) Calcium as a coagonist of inositol 1,4,5-trisphosphate-induced calcium release. *Science* 252:443–446
27. Iino M (1990) Biphasic Ca<sup>2+</sup> dependence of inositol 1,4,5-trisphosphate-induced Ca<sup>2+</sup> release in smooth muscle cells of the guinea pig taenia caeci. *J Gen Physiol* 95:1103–1122
28. Parys JB, Sernett SW, DeLisle S et al (1992) Isolation, characterization, and localization of the inositol 1,4,5-trisphosphate receptor protein in *Xenopus laevis* oocytes. *J Biol Chem* 267:18776–18782
29. Bezprozvanny I, Ehrlich BE (1993) ATP modulates the function of inositol 1,4,5-trisphosphate-gated channels at two sites. *Neuron* 10:1175–1184
30. Bootman MD, Taylor CW, Berridge MJ (1992) The thiol reagent, thimerosal, evokes Ca<sup>2+</sup> spikes in HeLa cells by sensitizing the inositol 1,4,5-trisphosphate receptor. *J Biol Chem* 267:25113–25119
31. Parys JB, Missiaen L, De Smedt H et al (1993) Bell-shaped activation of inositol-1,4,5-trisphosphate-induced Ca<sup>2+</sup> release by thimerosal in permeabilized A7r5 smooth-muscle cells. *Pflügers Arch* 424:516–522
32. Tovey SC, Dedos SG, Taylor EJA et al (2008) Selective coupling of type 6 adenylyl cyclase with type 2 IP<sub>3</sub> receptors mediates direct sensitization of IP<sub>3</sub> receptors by cAMP. *J Cell Biol* 183:297–311. <https://doi.org/10.1083/jcb.200803172>
33. Tovey SC, Dedos SG, Rahman T et al (2010) Regulation of inositol 1,4,5-trisphosphate receptors by cAMP independent of cAMP-dependent protein kinase. *J Biol Chem* 285:12979–12989. <https://doi.org/10.1074/jbc.M109.096016>
34. Konieczny V, Tovey SC, Mataragka S et al (2017) Cyclic AMP recruits a discrete intracellular Ca<sup>2+</sup> store by unmasking hypersensitive IP<sub>3</sub> receptors. *Cell Rep* 18:711–722. <https://doi.org/10.1016/j.celrep.2016.12.058>

35. Patterson RL, Boehning D, Snyder SH (2004) Inositol 1,4,5-trisphosphate receptors as signal integrators. *Annu Rev Biochem* 73:437–465
36. Vervloessem T, Yule DI, Bultynck G, Parys JB (2015) The type 2 inositol 1,4,5-trisphosphate receptor, emerging functions for an intriguing Ca<sup>2+</sup>-release channel. *Biochim Biophys Acta* 1853:1992–2005. <https://doi.org/10.1016/j.bbamcr.2014.12.006>
37. Iwai M, Michikawa T, Bosanac I et al (2007) Molecular basis of the isoform-specific ligand-binding affinity of inositol 1,4,5-trisphosphate receptors. *J Biol Chem* 282:12755–12764. <https://doi.org/10.1074/jbc.M609833200>
38. Miyakawa T, Maeda A, Yamazawa T et al (1999) Encoding of Ca<sup>2+</sup> signals by differential expression of IP<sub>3</sub> receptor subtypes. *EMBO J* 18:1303–1308. <https://doi.org/10.1093/emboj/18.5.1303>
39. Newton CL, Mignery GA, Südhof TC (1994) Co-expression in vertebrate tissues and cell lines of multiple inositol 1,4,5-trisphosphate (InsP<sub>3</sub>) receptors with distinct affinities for InsP<sub>3</sub>. *J Biol Chem* 269:28613–28619
40. Tu H, Wang Z, Nosyreva E et al (2005b) Functional characterization of mammalian inositol 1,4,5-trisphosphate receptor isoforms. *Biophys J* 88:1046–1055. <https://doi.org/10.1529/biophysj.104.049593>
41. Vanlingen S, Sipma H, De Smet P et al (2000) Ca<sup>2+</sup> and calmodulin differentially modulate myo-inositol 1,4, 5-trisphosphate (IP<sub>3</sub>)-binding to the recombinant ligand-binding domains of the various IP<sub>3</sub> receptor isoforms. *Biochem J* 346:275–280
42. Mak DO, McBride S, Foscett JK (2001) Regulation by Ca<sup>2+</sup> and inositol 1,4,5-trisphosphate (InsP<sub>3</sub>) of single recombinant type 3 InsP<sub>3</sub> receptor channels. Ca<sup>2+</sup> activation uniquely distinguishes types 1 and 3 InsP<sub>3</sub> receptors. *J Gen Physiol* 117:435–446
43. Mak D-OD, McBride SMJ, Petrenko NB, Foscett JK (2003) Novel regulation of calcium inhibition of the inositol 1,4,5-trisphosphate receptor calcium-release channel. *J Gen Physiol* 122:569–581. <https://doi.org/10.1085/jgp.200308808>
44. Tu H, Wang Z, Bezprozvanny I (2005a) Modulation of mammalian inositol 1,4,5-trisphosphate receptor isoforms by calcium: a role of calcium sensor region. *Biophys J* 88:1056–1069. <https://doi.org/10.1529/biophysj.104.049601>
45. Betzenhauser MJ, Wagner LE, Iwai M et al (2008) ATP modulation of Ca<sup>2+</sup> release by type-2 and type-3 inositol (1, 4, 5)-triphosphate receptors. Differing ATP sensitivities and molecular determinants of action. *J Biol Chem* 283:21579–21587. <https://doi.org/10.1074/jbc.M801680200>
46. Missiaen L, Parys JB, Sienaert I et al (1998) Functional properties of the type-3 InsP<sub>3</sub> receptor in 16HBE14o- bronchial mucosal cells. *J Biol Chem* 273:8983–8986
47. Maes K, Missiaen L, De Smet P et al (2000) Differential modulation of inositol 1,4,5-trisphosphate receptor type 1 and type 3 by ATP. *Cell Calcium* 27:257–267. <https://doi.org/10.1054/ceca.2000.0121>
48. Alzayady KJ, Wagner LE, Chandrasekhar R et al (2013) Functional inositol 1,4,5-trisphosphate receptors assembled from concatenated homo- and heteromeric subunits. *J Biol Chem* 288:29772–29784. <https://doi.org/10.1074/jbc.M113.502203>
49. Bultynck G, Szlufcik K, Kasri NN et al (2004) Thimerosal stimulates Ca<sup>2+</sup> flux through inositol 1,4,5-trisphosphate receptor type 1, but not type 3, via modulation of an isoform-specific Ca<sup>2+</sup>-dependent intramolecular interaction. *Biochem J* 381:87–96. <https://doi.org/10.1042/BJ20040072>
50. Bultynck G, Sienaert I, Parys JB et al (2003a) Pharmacology of inositol trisphosphate receptors. *Pflügers Arch* 445:629–642. <https://doi.org/10.1007/s00424-002-0971-1>
51. Saleem H, Tovey SC, Molinski TF, Taylor CW (2014) Interactions of antagonists with subtypes of inositol 1,4,5-trisphosphate (IP<sub>3</sub>) receptor. *Br J Pharmacol* 171:3298–3312. <https://doi.org/10.1111/bph.12685>
52. Swarbrick JM, Riley AM, Mills SJ, Potter BVL (2015) Designer small molecules to target calcium signalling. *Biochem Soc Trans* 43:417–425. <https://doi.org/10.1042/BST20140293>
53. Fredericks GJ, Hoffmann FW, Rose AH et al (2014) Stable expression and function of the inositol 1,4,5-triphosphate receptor requires palmitoylation by a DHHC6/selenoprotein K



- complex. *Proc Natl Acad Sci USA* 111:16478–16483. <https://doi.org/10.1073/pnas.1417176111>
54. Hamada K, Terauchi A, Nakamura K et al (2014) Aberrant calcium signaling by transglutaminase-mediated posttranslational modification of inositol 1,4,5-trisphosphate receptors. *Proc Natl Acad Sci USA* 111:E3966–E3975. <https://doi.org/10.1073/pnas.1409730111>
  55. Bultynck G, Vermassen E, Szlufcik K et al (2003b) Calcineurin and intracellular  $\text{Ca}^{2+}$ -release channels: regulation or association? *Biochem Biophys Res Commun* 311:1181–1193
  56. Vanderheyden V, Devogelaere B, Missiaen L et al (2009) Regulation of inositol 1,4,5-trisphosphate-induced  $\text{Ca}^{2+}$  release by reversible phosphorylation and dephosphorylation. *Biochim Biophys Acta* 1793:959–970. <https://doi.org/10.1016/j.bbamcr.2008.12.003>
  57. Prole DL, Taylor CW (2016) Inositol 1,4,5-trisphosphate receptors and their protein partners as signalling hubs. *J Physiol* 594:2849–2866. <https://doi.org/10.1113/JP271139>
  58. Ando H, Mizutani A, Kiefer H et al (2006) IRBIT suppresses  $\text{IP}_3$  receptor activity by competing with  $\text{IP}_3$  for the common binding site on the  $\text{IP}_3$  receptor. *Mol Cell* 22:795–806. <https://doi.org/10.1016/j.molcel.2006.05.017>
  59. Devogelaere B, Beullens M, Sammels E et al (2007) Protein phosphatase-1 is a novel regulator of the interaction between IRBIT and the inositol 1,4,5-trisphosphate receptor. *Biochem J* 407:303–311. <https://doi.org/10.1042/BJ20070361>
  60. Michikawa T, Hirota J, Kawano S et al (1999) Calmodulin mediates calcium-dependent inactivation of the cerebellar type 1 inositol 1,4,5-trisphosphate receptor. *Neuron* 23:799–808
  61. Missiaen L, Parys JB, Weidema AF et al (1999) The bell-shaped  $\text{Ca}^{2+}$  dependence of the inositol 1,4, 5-trisphosphate-induced  $\text{Ca}^{2+}$  release is modulated by  $\text{Ca}^{2+}$ /calmodulin. *J Biol Chem* 274:13748–13751
  62. Patel S, Morris SA, Adkins CE et al (1997)  $\text{Ca}^{2+}$ -independent inhibition of inositol trisphosphate receptors by calmodulin: redistribution of calmodulin as a possible means of regulating  $\text{Ca}^{2+}$  mobilization. *Proc Natl Acad Sci USA* 94:11627–11632
  63. Sienaert I, Nadif Kasri N, Vanlingen S et al (2002) Localization and function of a calmodulin-apocalmodulin-binding domain in the N-terminal part of the type 1 inositol 1,4,5-trisphosphate receptor. *Biochem J* 365:269–277. <https://doi.org/10.1042/BJ20020144>
  64. Yamada M, Miyawaki A, Saito K et al (1995) The calmodulin-binding domain in the mouse type 1 inositol 1,4,5-trisphosphate receptor. *Biochem J* 308:83–88
  65. Haynes LP, Tepikin AV, Burgoyne RD (2004) Calcium-binding protein 1 is an inhibitor of agonist-evoked, inositol 1,4,5-trisphosphate-mediated calcium signaling. *J Biol Chem* 279:547–555. <https://doi.org/10.1074/jbc.M309617200>
  66. Kasri NN, Holmes AM, Bultynck G et al (2004) Regulation of  $\text{InsP}_3$  receptor activity by neuronal  $\text{Ca}^{2+}$ -binding proteins. *EMBO J* 23:312–321. <https://doi.org/10.1038/sj.emboj.7600037>
  67. Yang J, McBride S, Mak D-OD et al (2002) Identification of a family of calcium sensors as protein ligands of inositol trisphosphate receptor  $\text{Ca}^{2+}$  release channels. *Proc Natl Acad Sci USA* 99:7711–7716. <https://doi.org/10.1073/pnas.102006299>
  68. Chen R, Valencia I, Zhong F et al (2004) Bcl-2 functionally interacts with inositol 1,4,5-trisphosphate receptors to regulate calcium release from the ER in response to inositol 1,4,5-trisphosphate. *J Cell Biol* 166:193–203. <https://doi.org/10.1083/jcb.200309146>
  69. Eckenrode EF, Yang J, Velmurugan GV et al (2010) Apoptosis protection by Mcl-1 and Bcl-2 modulation of inositol 1,4,5-trisphosphate receptor-dependent  $\text{Ca}^{2+}$  signaling. *J Biol Chem* 285:13678–13684. <https://doi.org/10.1074/jbc.M109.096040>
  70. Rong Y-P, Bultynck G, Aromolaran AS et al (2009) The BH4 domain of Bcl-2 inhibits ER calcium release and apoptosis by binding the regulatory and coupling domain of the  $\text{IP}_3$  receptor. *Proc Natl Acad Sci USA* 106:14397–14402. <https://doi.org/10.1073/pnas.0907555106>
  71. White C, Li C, Yang J et al (2005) The endoplasmic reticulum gateway to apoptosis by Bcl-X<sub>L</sub> modulation of the  $\text{InsP}_3\text{R}$ . *Nat Cell Biol* 7:1021–1028. <https://doi.org/10.1038/ncb1302>

72. Xu S, Xu Y, Chen L et al (2017) RCN1 suppresses ER stress-induced apoptosis via calcium homeostasis and PERK-CHOP signaling. *Oncogenesis* 6:e304. <https://doi.org/10.1038/oncsis.2017.6>
73. Decuypere J-P, Welkenhuyzen K, Luyten T et al (2011e) Ins(1,4,5)P<sub>3</sub> receptor-mediated Ca<sup>2+</sup> signaling and autophagy induction are interrelated. *Autophagy* 7:1472–1489
74. Vicencio JM, Ortiz C, Criollo A et al (2009) The inositol 1,4,5-trisphosphate receptor regulates autophagy through its interaction with Beclin 1. *Cell Death Differ* 16:1006–1017. <https://doi.org/10.1038/cdd.2009.34>
75. Higo T, Hattori M, Nakamura T et al (2005) Subtype-specific and ER luminal environment-dependent regulation of inositol 1,4,5-trisphosphate receptor type 1 by ERp44. *Cell* 120:85–98. <https://doi.org/10.1016/j.cell.2004.11.048>
76. Higo T, Hamada K, Hisatsune C et al (2010) Mechanism of ER stress-induced brain damage by IP<sub>3</sub> receptor. *Neuron* 68:865–878. <https://doi.org/10.1016/j.neuron.2010.11.010>
77. Hayashi T, Su TP (2001) Regulating ankyrin dynamics: Roles of sigma-1 receptors. *Proc Natl Acad Sci USA* 98:491–496. <https://doi.org/10.1073/pnas.021413698>
78. Boehning D, Patterson RL, Sedaghat L et al (2003) Cytochrome c binds to inositol (1,4,5) trisphosphate receptors, amplifying calcium-dependent apoptosis. *Nat Cell Biol* 5:1051–1061. <https://doi.org/10.1038/ncb1063>
79. Akl H, Bultynck G (2013) Altered Ca<sup>2+</sup> signaling in cancer cells: proto-oncogenes and tumor suppressors targeting IP<sub>3</sub> receptors. *Biochim Biophys Acta* 1835:180–193. <https://doi.org/10.1016/j.bbcan.2012.12.001>
80. Ivanova H, Kerkhofs M, La Rovere R, Bultynck G (2017) Endoplasmic reticulum–mitochondrial Ca<sup>2+</sup> fluxes underlying cancer cell survival. *Front Oncol* 7:Article 70. <https://doi.org/10.3389/fonc.2017.00070>
81. Decrock E, De Bock M, Wang N et al (2013) IP<sub>3</sub>, a small molecule with a powerful message. *Biochim Biophys Acta* 1833:1772–1786. <https://doi.org/10.1016/j.bbamcr.2012.12.016>
82. Distelhorst CW, Bootman MD (2011) Bcl-2 interaction with the inositol 1,4,5-trisphosphate receptor: role in Ca<sup>2+</sup> signaling and disease. *Cell Calcium* 50:234–241. <https://doi.org/10.1016/j.ceca.2011.05.011>
83. Ivanova H, Vervliet T, Missiaen L et al (2014) Inositol 1,4,5-trisphosphate receptor-isoform diversity in cell death and survival. *Biochim Biophys Acta* 1843:2164–2183. <https://doi.org/10.1016/j.bbamcr.2014.03.007>
84. Joseph SK, Hajnóczky G (2007) IP<sub>3</sub> receptors in cell survival and apoptosis: Ca<sup>2+</sup> release and beyond. *Apoptosis* 12:951–968. <https://doi.org/10.1007/s10495-007-0719-7>
85. La Rovere RML, Roest G, Bultynck G, Parys JB (2016) Intracellular Ca<sup>2+</sup> signaling and Ca<sup>2+</sup> microdomains in the control of cell survival, apoptosis and autophagy. *Cell Calcium* 60:74–87. <https://doi.org/10.1016/j.ceca.2016.04.005>
86. Smaili SS, Pereira GJS, Costa MM et al (2013) The role of calcium stores in apoptosis and autophagy. *Curr Mol Med* 13:252–265
87. Bootman MD, Chehab T, Bultynck G, et al Autophagy and calcium signalling: do we have a consensus? *Cell Calcium*. 2017; (submitted).
88. Ghislat G, Knecht E (2013) Ca<sup>2+</sup>-sensor proteins in the autophagic and endocytic traffic. *Curr Protein Pept Sci* 14:97–110
89. Parys JB, Decuypere J-P, Bultynck G (2012) Role of the inositol 1,4,5-trisphosphate receptor/Ca<sup>2+</sup>-release channel in autophagy. *Cell Commun Signal CCS* 10:17. <https://doi.org/10.1186/1478-811X-10-17>
90. Sun F, Xu X, Wang X, Zhang B (2016) Regulation of autophagy by Ca<sup>2+</sup>. *Tumour Biol* 37 (12):15467–15476. <https://doi.org/10.1007/s13277-016-5353-y>
91. Czabotar PE, Lessene G, Strasser A, Adams JM (2014) Control of apoptosis by the BCL-2 protein family: implications for physiology and therapy. *Nat Rev Mol Cell Biol* 15:49–63. <https://doi.org/10.1038/nrm3722>
92. Letai AG (2008) Diagnosing and exploiting cancer's addiction to blocks in apoptosis. *Nat Rev Cancer* 8:121–132. <https://doi.org/10.1038/nrc2297>

- 840 93. Tait SWG, Green DR (2013) Mitochondrial regulation of cell death. Cold Spring Harb  
841 Perspect Biol 5:a008706. <https://doi.org/10.1101/cshperspect.a008706>
- 842 94. Pinton P, Giorgi C, Siviero R et al (2008) Calcium and apoptosis: ER-mitochondria  $\text{Ca}^{2+}$   
843 transfer in the control of apoptosis. Oncogene 27:6407–6418. [https://doi.org/10.1038/onc.](https://doi.org/10.1038/onc.2008.308)  
844 2008.308
- 845 95. Hwang M-S, Schwall CT, Pazarentzos E et al (2014) Mitochondrial  $\text{Ca}^{2+}$  influx targets  
846 cardiolipin to disintegrate respiratory chain complex II for cell death induction. Cell Death  
847 Differ 21:1733–1745. <https://doi.org/10.1038/cdd.2014.84>
- 848 96. Bonora M, Bononi A, De Marchi E et al (2013) Role of the c subunit of the  $\text{F}_0$  ATP synthase in  
849 mitochondrial permeability transition. Cell Cycle 12:674–683. [https://doi.org/10.4161/cc.](https://doi.org/10.4161/cc.23599)  
850 23599
- 851 97. Giorgio V, von Stockum S, Antoniel M et al (2013) Dimers of mitochondrial ATP synthase  
852 form the permeability transition pore. Proc Natl Acad Sci USA 110:5887–5892. [https://doi.](https://doi.org/10.1073/pnas.1217823110)  
853 [org/10.1073/pnas.1217823110](https://doi.org/10.1073/pnas.1217823110)
- 854 98. Giorgio V, Guo L, Bassot C, et al Calcium and regulation of the mitochondrial permeability  
855 transition. Cell Calcium, 2017b: (submitted).
- 856 99. Giorgio V, Burchell V, Schiavone M et al (2017a)  $\text{Ca}^{2+}$  binding to F-ATP synthase  $\beta$  subunit  
857 triggers the mitochondrial permeability transition. EMBO Rep 18(7):1065–1076
- 858 100. Bender T, Martinou J-C (2013) Where killers meet--permeabilization of the outer mitochon-  
859 drial membrane during apoptosis. Cold Spring Harb Perspect Biol 5:a011106. [https://doi.org/](https://doi.org/10.1101/cshperspect.a011106)  
860 [10.1101/cshperspect.a011106](https://doi.org/10.1101/cshperspect.a011106)
- 861 101. Wang HG, Pathan N, Ethell IM et al (1999)  $\text{Ca}^{2+}$ -induced apoptosis through calcineurin  
862 dephosphorylation of BAD. Science 284:339–343
- 863 102. Sano R, Reed JC (2013) ER stress-induced cell death mechanisms. Biochim Biophys Acta  
864 1833:3460–3470. <https://doi.org/10.1016/j.bbamcr.2013.06.028>
- 865 103. Urrea H, Dufey E, Lisbona F et al (2013) When ER stress reaches a dead end. Biochim Biophys  
866 Acta 1833:3507–3517. <https://doi.org/10.1016/j.bbamcr.2013.07.024>
- 867 104. Ravikumar B, Sarkar S, Davies JE et al (2010) Regulation of mammalian autophagy in  
868 physiology and pathophysiology. Physiol Rev 90:1383–1435. [https://doi.org/10.1152/](https://doi.org/10.1152/physrev.00030.2009)  
869 [physrev.00030.2009](https://doi.org/10.1152/physrev.00030.2009)
- 870 105. Liu Y, Levine B (2015) Autosis and autophagic cell death: the dark side of autophagy. Cell  
871 Death Differ 22:367–376. <https://doi.org/10.1038/cdd.2014.143>
- 872 106. Liu Y, Shoji-Kawata S, Sumpter RM et al (2013) Autosis is a  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase-regulated form  
873 of cell death triggered by autophagy-inducing peptides, starvation, and hypoxia-ischemia.  
874 Proc Natl Acad Sci USA 110:20364–20371. <https://doi.org/10.1073/pnas.1319661110>
- 875 107. Bento CF, Renna M, Ghislat G et al (2016) Mammalian autophagy: how does it work? Annu  
876 Rev Biochem 85:685–713. <https://doi.org/10.1146/annurev-biochem-060815-014556>
- 877 108. Wesselborg S, Stork B (2015) Autophagy signal transduction by ATG proteins: from hierarchies  
878 to networks. Cell Mol Life Sci 72:4721–4757. <https://doi.org/10.1007/s00018-015-2034-8>
- 879 109. Hamasaki M, Furuta N, Matsuda A et al (2013) Autophagosomes form at ER-mitochondria  
880 contact sites. Nature 495:389–393. <https://doi.org/10.1038/nature11910>
- 881 110. Hayashi-Nishino M, Fujita N, Noda T et al (2009) A subdomain of the endoplasmic reticulum  
882 forms a cradle for autophagosome formation. Nat Cell Biol 11:1433–1437. [https://doi.org/10.](https://doi.org/10.1038/ncb1991)  
883 [1038/ncb1991](https://doi.org/10.1038/ncb1991)
- 884 111. Ylä-Anttila P, Vihinen H, Jokitalo E, Eskelinen E-L (2009) 3D tomography reveals connec-  
885 tions between the phagophore and endoplasmic reticulum. Autophagy 5:1180–1185
- 886 112. Roberts R, Ktistakis NT (2013) Omegasomes: PI3P platforms that manufacture autophagosomes.  
887 Essays Biochem 55:17–27. <https://doi.org/10.1042/bse0550017>
- 888 113. Proikas-Cezanne T, Takacs Z, Dönnies P, Kohlbacher O (2015) WIPI proteins: essential  
889 PtdIns3P effectors at the nascent autophagosome. J Cell Sci 128:207–217. [https://doi.org/10.](https://doi.org/10.1242/jcs.146258)  
890 [1242/jcs.146258](https://doi.org/10.1242/jcs.146258)
- 891 114. Decuyper J-P, Kindt D, Luyten T et al (2013) mTOR-controlled autophagy requires intra-  
892 cellular  $\text{Ca}^{2+}$  signaling. PLoS One 8:e61020. <https://doi.org/10.1371/journal.pone.0061020>



115. Grote-meier A, Alers S, Pfisterer SG et al (2010) AMPK-independent induction of autophagy by cytosolic Ca<sup>2+</sup> increase. *Cell Signal* 22:914–925. <https://doi.org/10.1016/j.cellsig.2010.01.015>
116. Høyer-Hansen M, Bastholm L, Szyaniarowski P et al (2007) Control of macroautophagy by calcium, calmodulin-dependent kinase kinase-beta, and Bcl-2. *Mol Cell* 25:193–205. <https://doi.org/10.1016/j.molcel.2006.12.009>
117. Luyten T, Welkenhuyzen K, Roest G et al (2017) Resveratrol-induced autophagy is dependent on IP<sub>3</sub>Rs and on cytosolic Ca<sup>2+</sup>. *Biochim Biophys Acta* 1864:947–956. <https://doi.org/10.1016/j.bbamcr.2017.02.013>
118. Messai Y, Noman MZ, Hasmim M et al (2014) ITPR1 protects renal cancer cells against natural killer cells by inducing autophagy. *Cancer Res* 74:6820–6832. <https://doi.org/10.1158/0008-5472.CAN-14-0303>
119. Engedal N, Torgersen ML, Guldvik IJ et al (2013) Modulation of intracellular calcium homeostasis blocks autophagosome formation. *Autophagy* 9:1475–1490. <https://doi.org/10.4161/auto.25900>
120. Gulati P, Gaspers LD, Dann SG et al (2008) Amino acids activate mTOR complex 1 via Ca<sup>2+</sup>/CaM signaling to hVps34. *Cell Metab* 7:456–465. <https://doi.org/10.1016/j.cmet.2008.03.002>
121. Khan MT, Joseph SK (2010) Role of inositol trisphosphate receptors in autophagy in DT40 cells. *J Biol Chem* 285:16912–16920. <https://doi.org/10.1074/jbc.M110.114207>
122. Sarkar S, Floto RA, Berger Z et al (2005) Lithium induces autophagy by inhibiting inositol monophosphatase. *J Cell Biol* 170:1101–1111. <https://doi.org/10.1083/jcb.200504035>
123. Cárdenas C, Foskett JK (2012) Mitochondrial Ca<sup>2+</sup> signals in autophagy. *Cell Calcium* 52:44–51. <https://doi.org/10.1016/j.ceca.2012.03.001>
124. Decuypere J-P, Bultynck G, Parys JB (2011a) A dual role for Ca<sup>2+</sup> in autophagy regulation. *Cell Calcium* 50:242–250. <https://doi.org/10.1016/j.ceca.2011.04.001>
125. East DA, Campanella M (2013) Ca<sup>2+</sup> in quality control: an unresolved riddle critical to autophagy and mitophagy. *Autophagy* 9:1710–1719. <https://doi.org/10.4161/auto.25367>
126. Rimessi A, Bonora M, Marchi S et al (2013) Perturbed mitochondrial Ca<sup>2+</sup> signals as causes or consequences of mitophagy induction. *Autophagy* 9:1677–1686. <https://doi.org/10.4161/auto.24795>
127. Vicencio JM, Lavandero S, Szabadkai G (2010) Ca<sup>2+</sup>, autophagy and protein degradation: thrown off balance in neurodegenerative disease. *Cell Calcium* 47:112–121. <https://doi.org/10.1016/j.ceca.2009.12.013>
128. Decuypere J-P, Monaco G, Bultynck G et al (2011b) The IP<sub>3</sub> receptor-mitochondria connection in apoptosis and autophagy. *Biochim Biophys Acta* 1813:1003–1013. <https://doi.org/10.1016/j.bbamcr.2010.11.023>
129. Rizzuto R, De Stefani D, Raffaello A, Mammucari C (2012) Mitochondria as sensors and regulators of calcium signalling. *Nat Rev Mol Cell Biol* 13:566–578. <https://doi.org/10.1038/nrm3412>
130. Giacomello M, Pellegrini L (2016) The coming of age of the mitochondria-ER contact: a matter of thickness. *Cell Death Differ* 23:1417–1427. <https://doi.org/10.1038/cdd.2016.52>
131. Danese A, Patergnani S, Bonora M et al (2017) Calcium regulates cell death in cancer: Roles of the mitochondria and mitochondria-associated membranes (MAMs). *Biochim Biophys Acta* 1858(8):615–627. <https://doi.org/10.1016/j.bbmbio.2017.01.003>
132. Giorgi C, Missiroli S, Patergnani S et al (2015) Mitochondria-associated membranes: composition, molecular mechanisms, and physiopathological implications. *Antioxid Redox Signal* 22:995–1019. <https://doi.org/10.1089/ars.2014.6223>
133. Herrera-Cruz MS, Simmen T (2017) Of yeast, mice and men: MAMs come in two flavors. *Biol Direct* 12:3. <https://doi.org/10.1186/s13062-017-0174-5>
134. Naon D, Scorrano L (2014) At the right distance: ER-mitochondria juxtaposition in cell life and death. *Biochim Biophys Acta* 1843:2184–2194. <https://doi.org/10.1016/j.bbamcr.2014.05.011>

135. van Vliet AR, Verfaillie T, Agostinis P (2014) New functions of mitochondria associated membranes in cellular signaling. *Biochim Biophys Acta* 1843:2253–2262. <https://doi.org/10.1016/j.bbamcr.2014.03.009>
136. Vervliet T, Clerix E, Seitaj B et al (2017) Modulation of  $\text{Ca}^{2+}$  signaling by anti-apoptotic B-cell lymphoma 2 proteins at the endoplasmic reticulum–mitochondrial interface. *Front Oncol* 7:Article 75. <https://doi.org/10.3389/fonc.2017.00075>
137. Csordás G, Renken C, Várnai P et al (2006) Structural and functional features and significance of the physical linkage between ER and mitochondria. *J Cell Biol* 174:915–921. <https://doi.org/10.1083/jcb.200604016>
138. Filadi R, Theurey P, Pizzo P (2017b) The endoplasmic reticulum-mitochondria coupling in health and disease: Molecules, functions and significance. *Cell Calcium* 62:1–15. <https://doi.org/10.1016/j.ceca.2017.01.003>
139. Paillusson S, Stoica R, Gómez-Suaga P et al (2016) There's something wrong with my MAM; the ER-mitochondria axis and neurodegenerative diseases. *Trends Neurosci* 39:146–157. <https://doi.org/10.1016/j.tins.2016.01.008>
140. Szabadkai G, Bianchi K, Várnai P et al (2006) Chaperone-mediated coupling of endoplasmic reticulum and mitochondrial  $\text{Ca}^{2+}$  channels. *J Cell Biol* 175:901–911. <https://doi.org/10.1083/jcb.200608073>
141. Csordás G, Thomas AP, Hajnóczky G (1999) Quasi-synaptic calcium signal transmission between endoplasmic reticulum and mitochondria. *EMBO J* 18:96–108. <https://doi.org/10.1093/emboj/18.1.96>
142. Csordás G, Várnai P, Golenár T et al (2010) Imaging interorganelle contacts and local calcium dynamics at the ER-mitochondrial interface. *Mol Cell* 39:121–132. <https://doi.org/10.1016/j.molcel.2010.06.029>
143. Giacomello M, Drago I, Bortolozzi M et al (2010)  $\text{Ca}^{2+}$  hot spots on the mitochondrial surface are generated by  $\text{Ca}^{2+}$  mobilization from stores, but not by activation of store-operated  $\text{Ca}^{2+}$  channels. *Mol Cell* 38:280–290. <https://doi.org/10.1016/j.molcel.2010.04.003>
144. Murgia M, Rizzuto R (2015) Molecular diversity and pleiotropic role of the mitochondrial calcium uniporter. *Cell Calcium* 58:11–17. <https://doi.org/10.1016/j.ceca.2014.11.001>
145. de Brito OM, Scorrano L (2008) Mitofusin 2 tethers endoplasmic reticulum to mitochondria. *Nature* 456:605–610. <https://doi.org/10.1038/nature07534>
146. Naon D, Zaninello M, Giacomello M et al (2016) Critical reappraisal confirms that Mitofusin 2 is an endoplasmic reticulum-mitochondria tether. *Proc Natl Acad Sci USA* 113:11249–11254. <https://doi.org/10.1073/pnas.1606786113>
147. Cerqua C, Anesti V, Pyakurel A et al (2010) Trichoplein/mitostatin regulates endoplasmic reticulum-mitochondria juxtaposition. *EMBO Rep* 11:854–860. <https://doi.org/10.1038/embor.2010.151>
148. Sugiura A, Nagashima S, Tokuyama T et al (2013) MITOL regulates endoplasmic reticulum-mitochondria contacts via Mitofusin2. *Mol Cell* 51:20–34. <https://doi.org/10.1016/j.molcel.2013.04.023>
149. Cosson P, Marchetti A, Ravazzola M, Orci L (2012) Mitofusin-2 independent juxtaposition of endoplasmic reticulum and mitochondria: an ultrastructural study. *PLoS One* 7:e46293. <https://doi.org/10.1371/journal.pone.0046293>
150. Filadi R, Greotti E, Turacchio G et al (2015) Mitofusin 2 ablation increases endoplasmic reticulum-mitochondria coupling. *Proc Natl Acad Sci USA* 112:E2174–E2181. <https://doi.org/10.1073/pnas.1504880112>
151. Leal NS, Schreiner B, Pinho CM et al (2016) Mitofusin-2 knockdown increases ER-mitochondria contact and decreases amyloid  $\beta$ -peptide production. *J Cell Mol Med* 20:1686–1695. <https://doi.org/10.1111/jcmm.12863>
152. Wang PTC, Garcin PO, Fu M et al (2015a) Distinct mechanisms controlling rough and smooth endoplasmic reticulum contacts with mitochondria. *J Cell Sci* 128:2759–2765. <https://doi.org/10.1242/jcs.171132>

153. Filadi R, Greotti E, Turacchio G et al (2017a) On the role of Mitofusin 2 in endoplasmic reticulum-mitochondria tethering. *Proc Natl Acad Sci USA* 114:E2266–E2267. <https://doi.org/10.1073/pnas.1616040114>
154. Naon D, Zaninello M, Giacomello M et al (2017) Reply to Filadi et al.: does Mitofusin 2 tether or separate endoplasmic reticulum and mitochondria? *Proc Natl Acad Sci USA* 114:E2268–E2269. <https://doi.org/10.1073/pnas.1618610114>
155. Walker AK, Atkin JD (2011) Stress signaling from the endoplasmic reticulum: a central player in the pathogenesis of amyotrophic lateral sclerosis. *IUBMB Life* 63:754–763. <https://doi.org/10.1002/iub.520>
156. De Vos KJ, Mórotz GM, Stoica R et al (2012) VAPB interacts with the mitochondrial protein PTPIP51 to regulate calcium homeostasis. *Hum Mol Genet* 21:1299–1311. <https://doi.org/10.1093/hmg/ddr559>
157. Nishimura AL, Mitne-Neto M, Silva HC et al (2004) A mutation in the vesicle-trafficking protein VAPB causes late-onset spinal muscular atrophy and amyotrophic lateral sclerosis. *Am J Hum Genet* 75:822–831
158. Iwasawa R, Mahul-Mellier A-L, Datler C et al (2011) Fis1 and Bap31 bridge the mitochondria-ER interface to establish a platform for apoptosis induction. *EMBO J* 30:556–568. <https://doi.org/10.1038/emboj.2010.346>
159. Breckenridge DG, Stojanovic M, Marcellus RC, Shore GC (2003) Caspase cleavage product of BAP31 induces mitochondrial fission through endoplasmic reticulum calcium signals, enhancing cytochrome c release to the cytosol. *J Cell Biol* 160:1115–1127. <https://doi.org/10.1083/jcb.200212059>
160. Simmen T, Aslan JE, Blagoveshchenskaya AD et al (2005) PACS-2 controls endoplasmic reticulum-mitochondria communication and Bid-mediated apoptosis. *EMBO J* 24:717–729. <https://doi.org/10.1038/sj.emboj.7600559>
161. Verfaillie T, Rubio N, Garg AD et al (2012) PERK is required at the ER-mitochondrial contact sites to convey apoptosis after ROS-based ER stress. *Cell Death Differ* 19:1880–1891. <https://doi.org/10.1038/cdd.2012.74>
162. Gellerich FN, Gizatullina Z, Gainutdinov T et al (2013) The control of brain mitochondrial energization by cytosolic calcium: the mitochondrial gas pedal. *IUBMB Life* 65:180–190. <https://doi.org/10.1002/iub.1131>
163. McCormack JG, Halestrap AP, Denton RM (1990) Role of calcium ions in regulation of mammalian intramitochondrial metabolism. *Physiol Rev* 70:391–425
164. Glancy B, Balaban RS (2012) Role of mitochondrial Ca<sup>2+</sup> in the regulation of cellular energetics. *Biochemistry (Mosc)* 51:2959–2973. <https://doi.org/10.1021/bi2018909>
165. Shanmughapriya S, Rajan S, Hoffman NE et al (2015) Ca<sup>2+</sup> signals regulate mitochondrial metabolism by stimulating CREB-mediated expression of the mitochondrial Ca<sup>2+</sup> uniporter gene MCU. *Sci Signal* 8:ra23. <https://doi.org/10.1126/scisignal.2005673>
166. Sugawara H, Kurosaki M, Takata M, Kurosaki T (1997) Genetic evidence for involvement of type 1, type 2 and type 3 inositol 1,4,5-trisphosphate receptors in signal transduction through the B-cell antigen receptor. *EMBO J* 16:3078–3088
167. Wen H, Xu WJ, Jin X et al (2015) The roles of IP<sub>3</sub> receptor in energy metabolic pathways and reactive oxygen species homeostasis revealed by metabolomic and biochemical studies. *Biochim Biophys Acta* 1853:2937–2944. <https://doi.org/10.1016/j.bbamcr.2015.07.020>
168. Cárdenas C, Miller RA, Smith I et al (2010) Essential regulation of cell bioenergetics by constitutive InsP<sub>3</sub> receptor Ca<sup>2+</sup> transfer to mitochondria. *Cell* 142:270–283. <https://doi.org/10.1016/j.cell.2010.06.007>
169. Gómez-Suaga P, Paillusson S, Stoica R et al (2017) The ER-mitochondria tethering complex VAPB-PTPIP51 regulates autophagy. *Curr Biol* 27:371–385. <https://doi.org/10.1016/j.cub.2016.12.038>
170. Bultynck G (2016) Onco-IP<sub>3</sub>Rs feed cancerous cravings for mitochondrial Ca<sup>2+</sup>. *Trends Biochem Sci* 41:390–393. <https://doi.org/10.1016/j.tibs.2016.03.006>

- 1048 171. Cárdenas C, Müller M, McNeal A et al (2016) Selective vulnerability of cancer cells by  
1049 inhibition of  $\text{Ca}^{2+}$  transfer from endoplasmic reticulum to mitochondria. *Cell Rep* 15:219–220.  
1050 <https://doi.org/10.1016/j.celrep.2016.03.045>
- 1051 172. Krols M, Bultynck G, Janssens S (2016) ER-Mitochondria contact sites: a new regulator of  
1052 cellular calcium flux comes into play. *J Cell Biol* 214:367–370. [https://doi.org/10.1083/jcb.](https://doi.org/10.1083/jcb.201607124)  
1053 [201607124](https://doi.org/10.1083/jcb.201607124)
- 1054 173. Raturi A, Gutiérrez T, Ortiz-Sandoval C et al (2016) TMX1 determines cancer cell metabolism  
1055 as a thiol-based modulator of ER-mitochondria  $\text{Ca}^{2+}$  flux. *J Cell Biol* 214:433–444. [https://doi.](https://doi.org/10.1083/jcb.201512077)  
1056 [org/10.1083/jcb.201512077](https://doi.org/10.1083/jcb.201512077)
- 1057 174. Granatiero V, Giorgio V, Cali T et al (2016) Reduced mitochondrial  $\text{Ca}^{2+}$  transients stimulate  
1058 autophagy in human fibroblasts carrying the 13514A>G mutation of the ND5 subunit of  
1059 NADH dehydrogenase. *Cell Death Differ* 23:231–241. <https://doi.org/10.1038/cdd.2015.84>
- 1060 175. MacVicar TDB, Mannack LVJC, Lees RM, Lane JD (2015) Targeted siRNA screens identify  
1061 ER-to-mitochondrial calcium exchange in autophagy and mitophagy responses in RPE1 cells.  
1062 *Int J Mol Sci* 16:13356–13380. <https://doi.org/10.3390/ijms160613356>
- 1063 176. Hayashi T, Su T-P (2007) Sigma-1 receptor chaperones at the ER-mitochondrion interface  
1064 regulate  $\text{Ca}^{2+}$  signaling and cell survival. *Cell* 131:596–610. [https://doi.org/10.1016/j.cell.](https://doi.org/10.1016/j.cell.2007.08.036)  
1065 [2007.08.036](https://doi.org/10.1016/j.cell.2007.08.036)
- 1066 177. Chan J, Yamazaki H, Ishiyama N et al (2010) Structural studies of inositol 1,4,5-trisphosphate  
1067 receptor: coupling ligand binding to channel gating. *J Biol Chem* 285:36092–36099. [https://doi.](https://doi.org/10.1074/jbc.M110.140160)  
1068 [org/10.1074/jbc.M110.140160](https://doi.org/10.1074/jbc.M110.140160)
- 1069 178. Williams A, Hayashi T, Wolozny D et al (2016) The non-apoptotic action of Bcl-xL:  
1070 regulating  $\text{Ca}^{2+}$  signaling and bioenergetics at the ER-mitochondrion interface. *J Bioenerg*  
1071 *Biomembr* 48:211–225. <https://doi.org/10.1007/s10863-016-9664-x>
- 1072 179. Yang J, Vais H, Gu W, Foscett JK (2016) Biphasic regulation of  $\text{InsP}_3$  receptor gating by dual  
1073  $\text{Ca}^{2+}$  release channel BH3-like domains mediates Bcl-XL control of cell viability. *Proc Natl*  
1074 *Acad Sci USA* 113:E1953–E1962. <https://doi.org/10.1073/pnas.1517935113>
- 1075 180. Sung PJ, Tsai FD, Vais H et al (2013) Phosphorylated K-Ras limits cell survival by blocking  
1076 Bcl-xL sensitization of inositol trisphosphate receptors. *Proc Natl Acad Sci USA*  
1077 110:20593–20598. <https://doi.org/10.1073/pnas.1306431110>
- 1078 181. Mendes CCP, Gomes DA, Thompson M et al (2005) The type III inositol 1,4,5-trisphosphate  
1079 receptor preferentially transmits apoptotic  $\text{Ca}^{2+}$  signals into mitochondria. *J Biol Chem*  
1080 280:40892–40900. <https://doi.org/10.1074/jbc.M506623200>
- 1081 182. De Stefani D, Bononi A, Romagnoli A et al (2012) VDAC1 selectively transfers apoptotic  $\text{Ca}^{2+}$   
1082 signals to mitochondria. *Cell Death Differ* 19:267–273. [https://doi.org/10.1038/cdd.2011.](https://doi.org/10.1038/cdd.2011.92)  
1083 [92](https://doi.org/10.1038/cdd.2011.92)
- 1084 183. Akl H, Monaco G, La Rovere R et al (2013)  $\text{IP}_3\text{R}2$  levels dictate the apoptotic sensitivity of  
1085 diffuse large B-cell lymphoma cells to an  $\text{IP}_3\text{R}$ -derived peptide targeting the BH4 domain of  
1086 Bcl-2. *Cell Death Dis* 4:e632. <https://doi.org/10.1038/cddis.2013.140>
- 1087 184. Jayaraman T, Marks AR (1997) T cells deficient in inositol 1,4,5-trisphosphate receptor are  
1088 resistant to apoptosis. *Mol Cell Biol* 17:3005–3012
- 1089 185. Csordás G, Hajnóczky G (2001) Sorting of calcium signals at the junctions of endoplasmic  
1090 reticulum and mitochondria. *Cell Calcium* 29:249–262. <https://doi.org/10.1054/ceca.2000.0191>
- 1091 186. Chami M, Oulès B, Szabadkai G et al (2008) Role of SERCA1 truncated isoform in the  
1092 proapoptotic calcium transfer from ER to mitochondria during ER stress. *Mol Cell*  
1093 32:641–651. <https://doi.org/10.1016/j.molcel.2008.11.014>
- 1094 187. Akl H, Vervloessem T, Kiviluoto S et al (2014) A dual role for the anti-apoptotic Bcl-2 protein  
1095 in cancer: mitochondria versus endoplasmic reticulum. *Biochim Biophys Acta*  
1096 1843:2240–2252. <https://doi.org/10.1016/j.bbamcr.2014.04.017>
- 1097 188. Greenberg EF, Lavik AR, Distelhorst CW (2014) Bcl-2 regulation of the inositol 1,4,5-  
1098 trisphosphate receptor and calcium signaling in normal and malignant lymphocytes: potential  
1099 new target for cancer treatment. *Biochim Biophys Acta* 1843:2205–2210. [https://doi.org/10.](https://doi.org/10.1016/j.bbamcr.2014.03.008)  
1100 [1016/j.bbamcr.2014.03.008](https://doi.org/10.1016/j.bbamcr.2014.03.008)

189. Lewis A, Hayashi T, Su T-P, Betenbaugh MJ (2014) Bcl-2 family in inter-organelle modulation of calcium signaling; roles in bioenergetics and cell survival. *J Bioenerg Biomembr* 46:1–15. <https://doi.org/10.1007/s10863-013-9527-7>
190. Parys JB (2014) The IP<sub>3</sub> receptor as a hub for Bcl-2 family proteins in cell death control and beyond. *Sci Signal* 7:pe4. <https://doi.org/10.1126/scisignal.2005093>
191. Vervliet T, Parys JB, Bultynck G (2016) Bcl-2 proteins and calcium signaling: complexity beneath the surface. *Oncogene* 35:5079–5092. <https://doi.org/10.1038/onc.2016.31>
192. Rong Y-P, Aromolaran AS, Bultynck G et al (2008) Targeting Bcl-2-IP<sub>3</sub> receptor interaction to reverse Bcl-2's inhibition of apoptotic calcium signals. *Mol Cell* 31:255–265. <https://doi.org/10.1016/j.molcel.2008.06.014>
193. Monaco G, Decrock E, Akl H et al (2012) Selective regulation of IP<sub>3</sub>-receptor-mediated Ca<sup>2+</sup> signaling and apoptosis by the BH4 domain of Bcl-2 versus Bcl-XL. *Cell Death Differ* 19:295–309. <https://doi.org/10.1038/cdd.2011.97>
194. Monaco G, Decrock E, Arbel N et al (2015) The BH4 domain of anti-apoptotic Bcl-XL, but not that of the related Bcl-2, limits the voltage-dependent anion channel 1 (VDAC1)-mediated transfer of pro-apoptotic Ca<sup>2+</sup> signals to mitochondria. *J Biol Chem* 290:9150–9161. <https://doi.org/10.1074/jbc.M114.622514>
195. Zhong F, Harr MW, Bultynck G et al (2011) Induction of Ca<sup>2+</sup>-driven apoptosis in chronic lymphocytic leukemia cells by peptide-mediated disruption of Bcl-2-IP<sub>3</sub> receptor interaction. *Blood* 117:2924–2934. <https://doi.org/10.1182/blood-2010-09-307405>
196. Lavik AR, Zhong F, Chang M-J et al (2015) A synthetic peptide targeting the BH4 domain of Bcl-2 induces apoptosis in multiple myeloma and follicular lymphoma cells alone or in combination with agents targeting the BH3-binding pocket of Bcl-2. *Oncotarget* 6:27388–27402. <https://doi.org/10.18632/oncotarget.4489>
197. Greenberg EF, McColl KS, Zhong F et al (2015) Synergistic killing of human small cell lung cancer cells by the Bcl-2-inositol 1,4,5-trisphosphate receptor disruptor BIRD-2 and the BH3-mimetic ABT-263. *Cell Death Dis* 6:e2034. <https://doi.org/10.1038/cddis.2015.355>
198. Khan MT, Wagner L, Yule DI et al (2006) Akt kinase phosphorylation of inositol 1,4,5-trisphosphate receptors. *J Biol Chem* 281:3731–3737. <https://doi.org/10.1074/jbc.M509262200>
199. Marchi S, Rimessi A, Giorgi C et al (2008) Akt kinase reducing endoplasmic reticulum Ca<sup>2+</sup> release protects cells from Ca<sup>2+</sup>-dependent apoptotic stimuli. *Biochem Biophys Res Commun* 375:501–505. <https://doi.org/10.1016/j.bbrc.2008.07.153>
200. Szado T, Vanderheyden V, Parys JB et al (2008) Phosphorylation of inositol 1,4,5-trisphosphate receptors by protein kinase B/Akt inhibits Ca<sup>2+</sup> release and apoptosis. *Proc Natl Acad Sci USA* 105:2427–2432. <https://doi.org/10.1073/pnas.0711324105>
201. Marchi S, Marinello M, Bononi A et al (2012) Selective modulation of subtype III IP<sub>3</sub>R by Akt regulates ER Ca<sup>2+</sup> release and apoptosis. *Cell Death Dis* 3:e304. <https://doi.org/10.1038/cddis.2012.45>
202. Bononi A, Bonora M, Marchi S et al (2013) Identification of PTEN at the ER and MAMs and its regulation of Ca<sup>2+</sup> signaling and apoptosis in a protein phosphatase-dependent manner. *Cell Death Differ* 20:1631–1643. <https://doi.org/10.1038/cdd.2013.77>
203. Kuchay S, Giorgi C, Simoneschi D et al (2017) PTEN counteracts FBXL2 to promote IP<sub>3</sub>R3- and Ca<sup>2+</sup>-mediated apoptosis limiting tumour growth. *Nature* 546:554–558. <https://doi.org/10.1038/nature22965>
204. Ilyin GP, Rialland M, Glaire D, Guguen-Guillouzo C (1999) Identification of a novel Skp2-like mammalian protein containing F-box and leucine-rich repeats. *FEBS Lett* 459:75–79
205. Chen BB, Glasser JR, Coon TA, Mallampalli RK (2011) FBXL2 is a ubiquitin E3 ligase subunit that triggers mitotic arrest. *Cell Cycle* 10:3487–3494. <https://doi.org/10.4161/cc.10.20.17742>
206. Chen BB, Glasser JR, Coon TA, Mallampalli RK (2012) F-box protein FBXL2 exerts human lung tumor suppressor-like activity by ubiquitin-mediated degradation of cyclin D3 resulting in cell cycle arrest. *Oncogene* 31:2566–2579. <https://doi.org/10.1038/onc.2011.432>



- 1154 207. Giorgi C, Ito K, Lin H-K et al (2010) PML regulates apoptosis at endoplasmic reticulum by  
1155 modulating calcium release. *Science* 330:1247–1251. [https://doi.org/10.1126/science.  
1189157](https://doi.org/10.1126/science.1189157)
- 1157 208. Paillard M, Tubbs E, Thiebaut P-A et al (2013) Depressing mitochondria-reticulum interac-  
1158 tions protects cardiomyocytes from lethal hypoxia-reoxygenation injury. *Circulation*  
1159 128:1555–1565. <https://doi.org/10.1161/CIRCULATIONAHA.113.001225>
- 1160 209. Gomez L, Thiebaut P-A, Paillard M et al (2016) The SR/ER-mitochondria calcium crosstalk is  
1161 regulated by GSK3 $\beta$  during reperfusion injury. *Cell Death Differ* 23:313–322. [https://doi.org/  
10.1038/cdd.2015.101](https://doi.org/10.1038/cdd.2015.101)
- 1163 210. Wiel C, Lallet-Daher H, Gitenay D et al (2014) Endoplasmic reticulum calcium release  
1164 through ITPR2 channels leads to mitochondrial calcium accumulation and senescence. *Nat*  
1165 *Commun* 5:3792. <https://doi.org/10.1038/ncomms4792>
- 1166 211. Morgan AJ (2016) Ca<sup>2+</sup> dialogue between acidic vesicles and ER. *Biochem Soc Trans*  
1167 44:546–553. <https://doi.org/10.1042/BST20150290>
- 1168 212. Morgan AJ, Platt FM, Lloyd-Evans E, Galione A (2011) Molecular mechanisms of  
1169 endolysosomal Ca<sup>2+</sup> signalling in health and disease. *Biochem J* 439:349–374. [https://doi.  
org/10.1042/BJ20110949](https://doi.org/10.1042/BJ20110949)
- 1171 213. Penny CJ, Kilpatrick BS, Eden ER, Patel S (2015) Coupling acidic organelles with the ER  
1172 through Ca<sup>2+</sup> microdomains at membrane contact sites. *Cell Calcium* 58:387–396. [https://doi.  
org/10.1016/j.ceca.2015.03.006](https://doi.org/10.1016/j.ceca.2015.03.006)
- 1174 214. Xu H, Ren D (2015) Lysosomal physiology. *Annu Rev Physiol* 77:57–80. [https://doi.org/10.  
1146/annurev-physiol-021014-071649](https://doi.org/10.1146/annurev-physiol-021014-071649)
- 1176 215. Melchionda M, Pittman JK, Mayor R, Patel S (2016) Ca<sup>2+</sup>/H<sup>+</sup> exchange by acidic organelles  
1177 regulates cell migration in vivo. *J Cell Biol* 212:803–813. [https://doi.org/10.1083/jcb.  
201510019](https://doi.org/10.1083/jcb.201510019)
- 1179 216. Patel S, Cai X (2015) Evolution of acidic Ca<sup>2+</sup> stores and their resident Ca<sup>2+</sup>-permeable  
1180 channels. *Cell Calcium* 57:222–230. <https://doi.org/10.1016/j.ceca.2014.12.005>
- 1181 217. Galione A (2015) A primer of NAADP-mediated Ca<sup>2+</sup> signalling: From sea urchin eggs to  
1182 mammalian cells. *Cell Calcium* 58:27–47. <https://doi.org/10.1016/j.ceca.2014.09.010>
- 1183 218. Morgan AJ, Davis LC, Ruas M, Galione A (2015) TPC: the NAADP discovery channel?  
1184 *Biochem Soc Trans* 43:384–389. <https://doi.org/10.1042/BST20140300>
- 1185 219. Patel S (2015) Function and dysfunction of two-pore channels. *Sci Signal* 8:re7. [https://doi.  
org/10.1126/scisignal.aab3314](https://doi.org/10.1126/scisignal.aab3314)
- 1187 220. Allbritton NL, Meyer T, Stryer L (1992) Range of messenger action of calcium ion and  
1188 inositol 1,4,5-trisphosphate. *Science* 258:1812–1815
- 1189 221. Sukumaran P, Schaar A, Sun Y, Singh BB (2016) Functional role of TRP channels in  
1190 modulating ER stress and autophagy. *Cell Calcium* 60:123–132. [https://doi.org/10.1016/j.  
ceca.2016.02.012](https://doi.org/10.1016/j.ceca.2016.02.012)
- 1192 222. Waller-Evans H, Lloyd-Evans E (2015) Regulation of TRPML1 function. *Biochem Soc Trans*  
1193 43:442–446. <https://doi.org/10.1042/BST20140311>
- 1194 223. Wang W, Zhang X, Gao Q, Xu H (2014) TRPML1: an ion channel in the lysosome. *Handb*  
1195 *Exp Pharmacol* 222:631–645. [https://doi.org/10.1007/978-3-642-54215-2\\_24](https://doi.org/10.1007/978-3-642-54215-2_24)
- 1196 224. Galione A (2011) NAADP receptors. *Cold Spring Harb Perspect Biol* 3:a004036. [https://doi.  
org/10.1101/cshperspect.a004036](https://doi.org/10.1101/cshperspect.a004036)
- 1198 225. Grimm C, Chen C-C, Wahl-Schott C, Biel M (2017) Two-pore channels: catalyzers of  
1199 endolysosomal transport and function. *Front Pharmacol* 8:45. [https://doi.org/10.3389/fphar.  
2017.00045](https://doi.org/10.3389/fphar.2017.00045)
- 1201 226. Kilpatrick BS, Eden ER, Schapira AH et al (2013) Direct mobilisation of lysosomal Ca<sup>2+</sup>  
1202 triggers complex Ca<sup>2+</sup> signals. *J Cell Sci* 126:60–66. <https://doi.org/10.1242/jcs.118836>
- 1203 227. Fameli N, Ogunbayo OA, van Breemen C, Evans AM (2014) Cytoplasmic nanojunctions  
1204 between lysosomes and sarcoplasmic reticulum are required for specific calcium signaling.  
1205 *F1000Research* 3:93. <https://doi.org/10.12688/f1000research.3720.1>


228. Lam AKM, Galione A (2013) The endoplasmic reticulum and junctional membrane commu- 1206  
nication during calcium signaling. *Biochim Biophys Acta* 1833:2542–2559. <https://doi.org/10.1016/j.bbamcr.2013.06.004> 1207
229. Raiborg C, Wenzel EM, Stenmark H (2015) ER-endosome contact sites: molecular composi- 1208  
tions and functions. *EMBO J* 34:1848–1858. <https://doi.org/10.15252/embj.201591481> 1209
230. Cancela JM, Churchill GC, Galione A (1999) Coordination of agonist-induced Ca<sup>2+</sup>-signalling 1210  
patterns by NAADP in pancreatic acinar cells. *Nature* 398:74–76. <https://doi.org/10.1038/18032> 1211
231. Churchill GC, Galione A (2001) NAADP induces Ca<sup>2+</sup> oscillations via a two-pool mechanism 1212  
by priming IP<sub>3</sub>- and cADPR-sensitive Ca<sup>2+</sup> stores. *EMBO J* 20:2666–2671. <https://doi.org/10.1093/emboj/20.11.2666> 1213
232. Gerasimenko JV, Charlesworth RM, Sherwood MW et al (2015) Both RyRs and TPCs are 1214  
required for NAADP-induced intracellular Ca<sup>2+</sup> release. *Cell Calcium* 58:237–245. <https://doi.org/10.1016/j.ceca.2015.05.005> 1215
233. Morgan AJ, Davis LC, Wagner SKTY et al (2013) Bidirectional Ca<sup>2+</sup> signaling occurs between 1216  
the endoplasmic reticulum and acidic organelles. *J Cell Biol* 200:789–805. <https://doi.org/10.1083/jcb.201204078> 1217
234. López Sanjurjo CI, Tovey SC, Taylor CW (2014) Rapid recycling of Ca<sup>2+</sup> between IP<sub>3</sub>- 1218  
sensitive stores and lysosomes. *PloS One* 9:e11275. <https://doi.org/10.1371/journal.pone.0111275> 1219
235. López-Sanjurjo CI, Tovey SC, Prole DL, Taylor CW (2013) Lysosomes shape Ins(1,4,5)P<sub>3</sub>- 1220  
evoked Ca<sup>2+</sup> signals by selectively sequestering Ca<sup>2+</sup> released from the endoplasmic reticu- 1221  
lum. *J Cell Sci* 126:289–300. <https://doi.org/10.1242/jcs.116103> 1222
236. Kilpatrick BS, Yates E, Grimm C et al (2016) Endo-lysosomal TRP mucolipin-1 channels 1223  
trigger global ER Ca<sup>2+</sup> release and Ca<sup>2+</sup> influx. *J Cell Sci* 129:3859–3867. <https://doi.org/10.1242/jcs.190322> 1224
237. Decuypere J-P, Parys JB, Bultynck G (2015) ITPRs/inositol 1,4,5-trisphosphate receptors in 1225  
autophagy: From enemy to ally. *Autophagy* 11:1944–1948. <https://doi.org/10.1080/15548627.2015.1083666> 1226
238. Penny CJ, Kilpatrick BS, Han JM et al (2014) A computational model of lysosome-ER Ca<sup>2+</sup> 1227  
microdomains. *J Cell Sci* 127:2934–2943. <https://doi.org/10.1242/jcs.149047> 1228
239. Raffaello A, Mammucari C, Gherardi G, Rizzuto R (2016) Calcium at the center of cell 1229  
signaling: Interplay between endoplasmic reticulum, mitochondria, and lysosomes. *Trends Biochem Sci* 41:1035–1049. <https://doi.org/10.1016/j.tibs.2016.09.001> 1230
240. Ronco V, Potenza DM, Denti F et al (2015) A novel Ca<sup>2+</sup>-mediated cross-talk between 1231  
endoplasmic reticulum and acidic organelles: implications for NAADP-dependent Ca<sup>2+</sup> sig- 1232  
nalling. *Cell Calcium* 57:89–100. <https://doi.org/10.1016/j.ceca.2015.01.001> 1233
241. Gómez-Suaga P, Luzón-Toro B, Churamani D et al (2012) Leucine-rich repeat kinase 2 reg- 1234  
ulates autophagy through a calcium-dependent pathway involving NAADP. *Hum Mol Genet* 21:511–525. <https://doi.org/10.1093/hmg/ddr481> 1235
242. Pereira GJS, Hirata H, Fimia GM et al (2011) Nicotinic acid adenine dinucleotide phosphate 1236  
(NAADP) regulates autophagy in cultured astrocytes. *J Biol Chem* 286:27875–27881. <https://doi.org/10.1074/jbc.C110.216580> 1237
243. Lin P-H, Duann P, Komazaki S et al (2015) Lysosomal two-pore channel subtype 2 (TPC2) 1238  
regulates skeletal muscle autophagic signaling. *J Biol Chem* 290:3377–3389. <https://doi.org/10.1074/jbc.M114.608471> 1239
244. Lu Y, Hao B-X, Graeff R et al (2013) Two pore channel 2 (TPC2) inhibits autophagosomal- 1240  
lysosomal fusion by alkalinizing lysosomal pH. *J Biol Chem* 288:24247–24263. <https://doi.org/10.1074/jbc.M113.484253> 1241
245. Wong C-O, Li R, Montell C, Venkatachalam K (2012) Drosophila TRPML is required for 1242  
TORC1 activation. *Curr Biol* 22:1616–1621. <https://doi.org/10.1016/j.cub.2012.06.055> 1243

- 1257 246. Wang W, Gao Q, Yang M et al (2015b) Up-regulation of lysosomal TRPML1 channels is  
 1258 essential for lysosomal adaptation to nutrient starvation. *Proc Natl Acad Sci USA* 112:  
 1259 E1373–E1381. <https://doi.org/10.1073/pnas.1419669112>
- 1260 247. Li X, Rydzewski N, Hider A et al (2016) A molecular mechanism to regulate lysosome  
 1261 motility for lysosome positioning and tubulation. *Nat Cell Biol* 18:404–417. [https://doi.org/10.](https://doi.org/10.1038/ncb3324)  
 1262 [1038/ncb3324](https://doi.org/10.1038/ncb3324)
- 1263 248. García-Rúa V, Feijóo-Bandín S, Rodríguez-Penas D et al (2016) Endolysosomal two-pore  
 1264 channels regulate autophagy in cardiomyocytes. *J Physiol* 594:3061–3077. [https://doi.org/10.](https://doi.org/10.1113/JP271332)  
 1265 [1113/JP271332](https://doi.org/10.1113/JP271332)
- 1266 249. Manzoni C (2017) The LRRK2-macroautophagy axis and its relevance to Parkinson's disease.  
 1267 *Biochem Soc Trans* 45:155–162. <https://doi.org/10.1042/BST20160265>
- 1268 250. Hockey LN, Kilpatrick BS, Eden ER et al (2015) Dysregulation of lysosomal morphology by  
 1269 pathogenic LRRK2 is corrected by TPC2 inhibition. *J Cell Sci* 128:232–238. [https://doi.org/](https://doi.org/10.1242/jcs.164152)  
 1270 [10.1242/jcs.164152](https://doi.org/10.1242/jcs.164152)
- 1271 251. Medina DL, Di Paola S, Peluso I et al (2015) Lysosomal calcium signalling regulates  
 1272 autophagy through calcineurin and TFEB. *Nat Cell Biol* 17:288–299. [https://doi.org/10.](https://doi.org/10.1038/ncb3114)  
 1273 [1038/ncb3114](https://doi.org/10.1038/ncb3114)
- 1274 252. Settembre C, Di Malta C, Polito VA et al (2011) TFEB links autophagy to lysosomal  
 1275 biogenesis. *Science* 332:1429–1433. <https://doi.org/10.1126/science.1204592>
- 1276 253. Medina DL, Fraldi A, Bouche V et al (2011) Transcriptional activation of lysosomal exocy-  
 1277 tosis promotes cellular clearance. *Dev Cell* 21:421–430. [https://doi.org/10.1016/j.devcel.2011.](https://doi.org/10.1016/j.devcel.2011.07.016)  
 1278 [07.016](https://doi.org/10.1016/j.devcel.2011.07.016)
- 1279 254. Sardiello M, Palmieri M, di Ronza A et al (2009) A gene network regulating lysosomal  
 1280 biogenesis and function. *Science* 325:473–477. <https://doi.org/10.1126/science.1174447>



# Author Query

Chapter No.: 7	427856_1_En
----------------	-------------

Query Refs.	Details Required	Author's response
AU1	Please check the footnote, and correct if necessary.	

Uncorrected Proof